arXiv:1304.6271v1 [math.PR] 23 Apr 2013

Isotropic Markov semigroups on ultra-metric spaces

Alexander Bendikov* Alexander Grigor'yan† Christophe Pittet‡
Wolfgang Woess§

April 22, 2013

Abstract

Let (X, d) be a locally compact separable ultra-metric space. Given a reference measure μ on X and a step length distribution σ on $[0\,,\,\infty)$, we construct a symmetric Markov semigroup $\{P^t\}_{t\geq 0}$ acting in $L^2(X, \mu)$. Let $\{\mathcal{X}_t\}$ be the corresponding Markov process. We obtain upper and lower bounds of its transition density and its Green function, give a transience criterion, estimate its moments and describe the Markov generator $\mathcal{L}$ and its spectrum, which is pure point. In the particular case when X is the field of p-adic numbers, our construction recovers the Vladimirov Laplacian which is closely related to the concept of p-adic Quantum Mechanics. Even in this well established setting our results are new. We also elaborate the relation between our processes and Kigami's jump processes on the boundary of a tree which are induced by a random walk. In conclusion, we explain the fractional derivative on the p-adic integers and the corresponding random walk on the associated tree.

Contents

1 Introduction **2**

2 Heat semigroup and heat kernel **9**

3 Spectral Distribution Function **12**

4 Basic estimates of the Heat Kernel **16**

5 Green function and transience **21**

6 Moments of the Markov process **25**

7 The Markov generator and its spectrum **30**

8 The p-adic fractional derivative **40**

9 Vladimirov's Laplacian **45**

10 Random walks on a tree and jump processes on its boundary **49**

11 Duality of random walks on trees and isotropic processes on their boundaries **58**

12 Fractional derivative on p-adic integers and the associated random walk **65**

---

[*] Supported by the Polish Government Scientific Research Fund, Grant 2012/05/B/ST 1/00613

[†] Supported by SFB 701 of German Research Council

[‡] Supported by the CNRS

[§] Supported by Austrian Science Fund projects FWF W1230-N13 and FWF P24028-N18

1 Introduction

In this paper, we introduce and study a class of symmetric Markov processes defined on an ultra-metric space (X, d). Recall that a metric d is called an *ultra-metric* if it satisfies the ultra-metric inequality

$$d(x, y) \leq \max\{d(x, z), d(z, y)\}, \tag{1.1}$$

that is obviously stronger than the usual triangle inequality.

The ultra-metric property (1.1) implies that the balls in an ultra-metric space (X, d) look very differently from the classical Euclidean balls. Any two closed ultra-metric balls of the same radius are either disjoint or identical. The minimal radius of a ball coincides with its diameter. It follows that the collection of all balls of the same radius r forms a partition of X. This can be used to characterize the distance $d(x, y)$ between points $x, y \in X$ as follows: $d(x, y)$ equals to the minimal value of r such that both x and y belong to the same ball of radius r. This property in turn implies that to define an ultra-metric d on X one has to prescribe to each ball B the value $d(B)$ of its diameter; see §10.

One of the best known examples of an ultra-metric space is the field $\mathbb{Q}_p$ of p-adic numbers endowed with the p-adic norm $\|x\|_p$ and the $p-$adic ultra-metric $d(x, y) = \|x - y\|_p$. Moreover, for any integer $n \geq 1$, the p-adic n-space $\mathbb{Q}_p^n = \mathbb{Q}_p \times ... \times \mathbb{Q}_p$ is also an ultra-metric space with the ultra-metric $d_n(x, y)$ defined as

$$d_n(x, y) = \max\{d(x_1, y_1), ..., d(x_n, y_n)\}.$$

All ultra-metric spaces considered in this paper are assumed to satisfy two natural conditions:

(i) (X, d) is complete and separable.

(ii) Every closed ball $B_r(x) = \{y \in X : d(x, y) \leq r\}$ is compact.

If in addition (X, d) satisfies the third condition:

(iii) The group of isometries of (X, d) acts transitively on X,

then (X, d) is in fact a locally compact Abelian group, that can be identified with any Abelian subgroup of its group of isometries that acts transitively on (X, d). For example, this is the case for $\mathbb{Q}_p^n$. Observe that this identification is not unique (!). With an eye on typical applications, we pay specific attention to the following two basic in the course of study cases:

Case 1.1 (X, d) *is discrete and infinite.*

Case 1.2 (X, d) *is perfect, that is, it contains no isolated point.*

Various constructions of Markov processes on non-compact perfect ultra-metric spaces (X, d) satisfying the hypotheses (i)–(iii) have been developed by DEL MUTO AND FIGÁ-TALAMANCA [28], [29], ALBEVERIO AND KARWOWSKI [1], ALBEVERIO AND ZHAO [2] and KOCHUBEI [26]. They fixed any identification of (X, d) with a locally compact Abelian group (in fact, $\mathbb{Q}_p$) and studied X-valued infinite divisible random variables and processes by using tools of Fourier analysis; for general references, see HEWITT AND ROSS [21], TAIBLESON [40] or SALOFF-COSTE [36]. Condition (iii) has later been relaxed but still all constructions relied crucially on an identification of (X, d) with some metric space of sequences. See also PEARSON AND BELLISSARD [31], and KIGAMI [24], [25], where (X, d) is the Cantor set, resp. the Cantor set minus one point. We shall come back to Kigami's work in §10.

An entirely different approach was developed by VLADIMIROV, VOLOVICH AND ZELENOV [44]. They were concerned with p-adic analysis (Bruhat distributions, Fourier transform etc.) related to the concept of p-adic Quantum Mechanics and introduced a class of pseudo-differential operators on $\mathbb{Q}_p$ and partially on $\mathbb{Q}_p^n$. In particular, they considered the p-adic Laplacian defined on $\mathbb{Q}_p^3$ and studied the corresponding p-adic Schrödinger equation. Among other results, they explicitly computed (as series expansions) certain heat kernels as well as the Green function of the p-adic Laplacian.

Ultra-metric spaces (X, d) that satisfy conditions (i)–(ii) and which are *discrete* were treated by Bendikov, Grigor'yan and Pittet [5], the direct forerunner of the present work. Among the examples of such spaces we mention the class of *locally finite* groups: a countable group G is locally finite if any of its finite subsets generates a finite subgroup. Every locally finite group G is the union of an increasing sequence of finite subgroups $\{G_n\}$. An ultra-metric d in G can be defined as follows: $d(x, y)$ is the minimal value of n such that x and y belong to a common coset of G_n.

Let us list a few well known examples of locally finite groups.

- $\mathbb{Z}(p) \oplus \mathbb{Z}(p) \oplus \cdots = \varinjlim \underbrace{\mathbb{Z}(p) \oplus \cdots \oplus \ \mathbb{Z}(p)}_{n \text{ times}}$ – the infinite sum of the cyclic groups $\mathbb{Z}(p)$;
- $S_\infty := \varinjlim S_n$ – the infinite symmetric group;
- $SL_\infty(p) := \varinjlim SL_n(p)$ - the infinite special linear group over the finite field $\mathbb{Z}(p)$.

Here are further interesting examples of ultrametric spaces.

- Boundaries, resp. punctured boundaries of locally finite trees - these are the generic examples, since every locally compact ultrametric space can be realized in this way. (Compare with §10 below.)
- Profinite groups, that is, groups which are inverse limits of finite groups. They are compact and totally disconnected; see Ribes and Zalesskii [35]. If such a group G has a first-countable topology then that topology is generated by an ultra-metric: first-countability amounts to the existence of a decreasing sequence of open normal subgroups N_k with trivial intersection. If $x, y \in G$ are distinct then there is a minimal k such that $xN_k \neq yN_k$, and $d(x, y) = 2^{-k}$ defines a suitable ultra-metric.[1]
- For any positive integer N, the collection of all isomorphism classes of finite or infinite connected, rooted graphs with vertex degrees bounded by N is well-known to be a compact ultrametric space. An ultrametric is given as follows: the distance between two non-isomorphic graphs (resp. representatives) is $1/r$, where r is the largest integer such that the balls of radius r around the root (with respect to the discrete graph metric) are isomorphic.
- A closed subspace of the last one is given by the collection of all finite or infinite rooted trees with vertex degrees bounded by N.

We briefly return to our discussion of locally finite groups. Since they are not finitely generated, the basic notions of geometric group theory such as the word metric, volume growth, isoperimetric inequalities, etc. (cf. e.g. Gromov [20]), do not have their counter-part in this setting. Hence one cannot apply directly the well-developed methods to study random walks on finitely generated groups provided by Varopoulos, Saloff-Coste and Coulhon [41], Coulhon, Grigor'yan and Pittet [14], Woess [46], Pittet and Saloff-Coste [34], [32],

---

[1] The 4th author thanks Wolfgang Herfort for a brief outline.

[33], Saloff-Coste [37], and others. Additional arguments are needed. The notion of an ultra-metric can be used instead of the word metric in this setting. See [5] and as well as Bendikov, Bobikau and Pittet [4], [3].

Selecting a set of generators for each subgroup G_n , one defines thereby a random walk on G_n, that is, a Markov kernel on G_n. Taking a convex combination of the Markov kernels across all G_n, one obtains a Markov kernel on G that determines a random walk on $G = \varinjlim G_n$. Such random walks have been studied by Darling and Erdös [15], Kesten and Spitzer [23], Flatto and Pitt [18], Fereig and Molchanov [17], Kasymdzhanova [22], Cartwright [12], Lawler [27], Brofferio and Woess [10]. In particular, [27] has a remarkable general criterion of recurrence of such random walks. The main novelty here is that many of those results are subsumed by our approach via ultra-metrics: the purpose of this paper is to develop tools to analyze a class of very natural and simply defined Markov processes on ultra-metric spaces (discrete and non-discrete as well) without assuming any group structure, but only under the hypotheses (*i*) and (*ii*). We do not need to use any identification of the ultra-metric space (X, d) with a set of sequences. In particular, this makes our arguments of completely geometric-analytic nature and allows us to bring into consideration an arbitrary Radon measure μ on X (instead of the Haar measure in the case of groups), that will be used as a speed measure for a Markov process. This approach has certain historical reminiscences: the theory of elliptic operators in $\mathbb{R}^n$ can be developed first in the case of constant coefficients by using the Fourier transform, whereas the case of variable coefficients requires more advanced tools. The measure μ in our case (see below) will play the role of a variable coefficient.

The aforementioned construction of random walks on locally finite groups can be easily generalized to an arbitrary ultra-metric measure space (X, d, μ) : instead of a sequence of subgroups $\{G_n\}_{n=1}^{\infty}$ one uses the family of balls $\{B_r(x)\}_{r>0, x\in X}$, and in each ball one has to choose a Markov kernel. The latter will be chosen as the μ-uniform distribution in each ball.

Here is our construction.

In addition to the measure μ on X, we fix a probability measure σ on the half-line $[0\,,\,\infty)$ with $\sigma(\{0\}) = 0$ and define the following Markov operators, where $r \geq 0$ and $f \in L^{\infty}(X, \mu)$.

$$\mathrm{Q}_r f(x) = \frac{1}{\mu(B_r(x))} \int_{B_r(x)} f\, d\mu, \tag{1.2}$$

$$Pf(x) = \int_0^{\infty} \mathrm{Q}_r f(x)\, d\sigma(r). \tag{1.3}$$

We associate with these operators two Markov processes.

1. A discrete time Markov chain $\{\mathcal{X}_n\}$, $n = 0, 1, ...$ with the following transition rule: first, a radius r is chosen at random according to the probability σ. Then $\mathcal{X}_{n+1}$ is μ-uniformly distributed in $B_r(\mathcal{X}_n)$.

2. A continuous time Markov process $\{\mathcal{X}_t\}_{t\geq 0}$ associated with the semigroup $\left\{P^t\right\}_{t\geq 0}$, where the operators P^t are given as

$$P^t f(x) = \int_0^{\infty} \mathrm{Q}_r f(x)\, d\sigma^t(r)\,, \qquad P^0 = id. \tag{1.4}$$

Here, σ^t is the probability measure on the half-line with $\sigma^t[0\,,\,r) = \big(\sigma[0\,,\,r)\big)^t$.

It follows that the discrete time Markov chain coincides with the restriction of the continuous time Markov process $\{\mathcal{X}_t\}$ to integer values of t, which allows us to study both simultaneously. (We will mostly consider continuous time.) It is suggestive to refer to those two processes as an *isotropic random walk,* resp. *isotropic jump process.* Each of them depends on three data: the ultra-metric d, the speed measure μ and the distance distribution σ. Thus, we call this the (d, μ, σ)*-process.* There is some degree of freedom. For example, we may modify the ultra-metric d and require that σ is a specific measure.

Definition 1.3 The *standard process* associated with μ and d is the one where $\sigma = \sigma_*$ is the "inverse exponential distribution" whose distribution function is

$$\sigma_* ([0\,,\, r)) = e^{-1/r}\,, \quad r > 0\,. \tag{1.5}$$

The *intrinsic ultra-metric* associated with $\{P^t\}$ on X is given by

$$\frac{1}{d_*(x,y)} = \log \frac{1}{\sigma\big[0\,,\, d(x,y)\big)}. \tag{1.6}$$

d_* induces the same collection of balls as d, and our process becomes the standard (d_*, μ)-process, as one easily verifies by the corresponding substitution in the integral appearing in (1.3), resp. (1.4).

An isotropic random walk on an ultra-metric measure space has a unique feature: its Markov operator P admits a spectral resolution in $L^2\,(X, \mu)$ where the spectral projectors are again Markov operators. This property brings up a new insight, new technical possibilities, and a new type of results, that have no analogue in other commonly used settings. For example, in our setting, the composition of any increasing function with the generator of the Markov semigroup is again a Markov generator. This property is in striking contrast with the classical result of Bochner about general Markov generators, where one needs to compose with a Bernstein function. In the context of random walks on groups, compare with BENDIKOV AND SALOFF-COSTE [7] as well as [6].

In the present study the following issues will be specifically dealt with.

(A) Explicit construction of the heat kernel (=transition probabilities) and the spectral distribution function (in §2 and §3).

(B) Estimates of the heat kernel and the Green function (in §4 and §5).

(C) In particular, criteria of recurrence and transience, including an analogue of Lawler's criterion without assuming any group structure (in §5).

(D) Estimates of escape rates (in §6).

(E) A precise description of the infinitesimal generator (= minus the "Laplacian") and its spectral properties (in §7).

(F) In particular, subordination techniques (also in §7): as it should be clear from the above remark, this will look entirely different from the classical setting.

Besides several examples that accompany the elaboration of our theory, we clarify with particular emphasis how that theory is related with two seemingly different analytic, resp. stochastic topics:

(G) The p-adic Laplace operator of Vladimirov (as well as related operators) embeds in our theory as the generator of one of our isotropic jump processes on $\mathbb{Q}_p$. (See §8, §9 and §12)

(H) The processes on the boundary of a tree that arise via "harmonic transform" from random walks on a tree, and that were recently examined by Kigami, embed naturally in our setting. (See §10, §11 and §12)

Let us return to a slightly more detailed outline of the analytic body of this work.

One of our aims is to obtain estimates of the transition density $p\,(t,x,y)$ of the process $\{\mathcal{X}_t\}$ (that is, the integral kernel of P^t) via the probability measure σ and the intrinsic volume function

$$V(x,r)=\mu\big(B_r^*(x)\big), \tag{1.7}$$

where $B_r^*(x)$ is the closed r-ball in the intrinsic metric d_*.

As indicated above, our approach is based upon the observation that the building blocks of the operator P, namely, the averaging operators Q_r of (1.2), are orthogonal projectors in $L^2(X,\mu)$. This very specific property is a consequence of the ultra-metric inequality (1.1). The range of Q_r coincides with the linear subspace $\mathcal{V}_r$ of $L^2(X,\mu)$ of all functions that are constant on each ball of radius r. Moreover, the spaces $\mathcal{V}_r$ are decreasing in r, so that the family $\{\mathrm{Q}_r\}$ is (up to reparametrization) a spectral resolution of the identity. Hence, the ultra-metric property leads to the spectral resolution (E_τ) of P as a self-adjoint operator in L^2, where the spectral projectors are also Markov operators. This enables us to engage at an early stage the methods of spectral theory and functional calculus.

Many of our results are stated in terms of the spectral distribution function N defined as

$$N(x,\tau)=1/V(x,1/\tau).$$

In the particular case of a discrete space X, we have $N(x,\tau)=(E_\tau \mathbf{1}_{\{x\}},\mathbf{1}_{\{x\}})$. If the group of isometries of (X,d) acts transitively on X, then $N(x,\tau)$ does not depend on x and coincides with the von Neumann trace of the orthoprojector E_τ. In general, $N(x,\tau)$ is a non-decreasing staircase function on $[0,\infty)$ whose behavior as $\tau\to 0$ is intimately related to the behavior of the heat kernel $p\,(t,x,y)$ as $t\to\infty$. Indeed, we have the following explicit identities

$$p(t,x,y)=t\int_0^{1/d_*(x,y)} N(x,\tau)\,\exp(-\tau t)\,d\tau, \tag{1.8}$$

$$p(t,x,x)=\int_0^{\infty}\exp(-\tau t)\,dN(x,\tau) \tag{1.9}$$

(Lebesgue-Stieltjes integral with respect to $\tau\mapsto N(x,\tau)$).

Fixing x and writing $N(\tau)=N(x,\tau)$, the equation (1.8) shows that

$$p(t,x,y)=\widehat{N}\left(t,\frac{1}{d_*(x,y)}\right),$$

where

$$\widehat{N}(t,r)=t\int_0^{r} N(\tau)\,\exp(-\tau t)\,d\tau$$

is the Laplace-type transform of the function $N(\tau)$. Our heat kernel estimates are based on the study of the function $\widehat{N}(t,r)$ as a functional of N. For example,

$$p\,(t,x,y)\simeq\frac{t}{t+d_*(x,y)}\,N\left(x,\frac{1}{t+d_*(x,y)}\right)$$

if and only if the function $N(\tau)$ satisfies the doubling property. (For the latter, see Definition 4.2 below. By $\simeq$, we indicate that the quotient of left and right hand side is bounded above and below by positive constants.) In particular, if $N\,(x,\tau)\simeq\tau^{\alpha}$ then

$$p(t,x,y)\simeq\frac{t}{\big(t^2+d_*(x,y)^2\big)^{\frac{1+\alpha}{2}}},$$

that is, $p\,(t,x,y)$ behaves like the Cauchy distribution in "α-dimensional" space. This example is closely related to studying p-adic fractional derivatives of order α.

The Green function estimates, as well as conditions for transience of the Markov process $\{\mathcal{X}_t\}$, are based on the fact that the Green function (more generally, the λ-Green function) is a Stieltjes-type transform of the spectral distribution function N. Using general properties of the Stieltjes transform, we obtain various results. For instance, the process is transient if and only if

$$\int_0 N(x,\tau)\,\frac{d\tau}{\tau^2} < \infty.$$

In particular, when $N(x,\tau) \simeq \tau^{\alpha}$ as $\tau \to 0$, transience is equivalent to $\alpha > 1$.

The identity (1.8) allows us to estimate the moments

$$M_\gamma(x,t) = \mathbb{E}_x(d_*(x,\mathcal{X}_t)^\gamma)$$

of the process $\{\mathcal{X}_t\}$. Indeed, we have the following explicit identity

$$M_\gamma(x,t) = \int_0^\infty \mathcal{R}_\gamma(x,\frac{t}{\tau})\,e^{-\tau}\,d\tau,$$

where

$$\mathcal{R}_\gamma(x,\tau) = \frac{1}{V(x,\tau)}\int_{(0,\tau]} s^\gamma\,dV(x,s)$$

(Lebesgue-Stieltjes integral with respect to the intrinsic volume measure $s \mapsto V(x,s)$).

In particular, we obtain the following general results:

(1) Assume that (X,d) is non-compact, has no isolated points, and that $0<\gamma<1$. Then

$$M_\gamma(x,t) \le \frac{t^\gamma}{1-\gamma}.$$

Thus in general the moment of order γ is finite for all $0<\gamma<1$. If $V(x,r)$ satisfies the reverse doubling condition – see Definition 4.9 below – then there is a matching lower bound. In this case the γ-moment is finite if and only if $0<\gamma<1$.

(2) Assume that (X,d) is discrete and infinite, and that $0<\gamma<1$. Then

$$M_\gamma(x,t) \le \frac{c}{1-\gamma}\min\{t,t^\gamma\}$$

for some $c>0$. If $V(x,r)$ satisfies the reverse doubling condition then there is a matching lower bound. In this case the γ-moment is finite if and only if $0<\gamma<1$.

(3) Assume that (X,d) is compact and perfect. then

$$M_\gamma(x,t) \le \begin{cases} C\,t, & \text{if} \quad \gamma>1,\\ C\,t\,(\log\frac{1}{t}+1), & \text{if} \quad \gamma=1,\\ C\,t^\gamma, & \text{if} \quad 0<\gamma<1,\end{cases}$$

for some $C>0$ and all $0<t\le 1$. If $V(x,r)$ satisfies the reverse doubling condition then there is again a matching lower bound.

If $-\mathcal{L}$ is the infinitesimal generator of the (d,μ,σ)-process then in our terminology, $\mathcal{L}$ is the *Laplacian.* We mention the following property: for any continuous, strictly increasing function $\varphi : [0\,,\,\infty) \to [0\,,\,\infty)$ such that $\varphi(0)=0$, the operator $\phi(\mathcal{L})$ is also a Laplacian of an analogously constructed Markov semigroup. In particular, $\mathcal{L}^\alpha$ is a Laplacian for any $\alpha>0$. Recall for comparison that, for a general symmetric Markov generator $-\mathcal{L}$, the operator $-\mathcal{L}^\alpha$ generates a Markov semigroup only for $0<\alpha\le 1$. Three further remarkable properties of $\mathcal{L}$ are:

- $\mathcal{L}$ extends to $L^p(X,\mu)$, and its spectrum in that space is the same for all $p \in [1\,,\,+\infty)$.

- Assume that (X, d) is non-compact. Let $M \subseteq [0\,,\,\infty)$ be a closed set that accumulates at 0 and such that M is unbounded if X contains at least one non-isolated point. Then there exist a proper ultra-metric d' on X which generates the same topology and an isotropic jump process on (Xd') whose associated Laplacian $\mathcal{L}$ has M as its spectrum.

- $\mathcal{L}$ has the strong Liouville property, that is, any non-negative $\mathcal{L}$-harmonic function must be constant.

We are able to apply our results to locally finite groups, but with arbitrary reference measure μ instead of the Haar (=counting) measure. Some of the above mentioned questions are particularly sensitive to the choice of the speed measure, for example, the heat kernel and Green function estimates. The spectrum of the Laplacian and escape rate bounds do not depend on the speed measure μ. These quantities depend strongly on the choice of the ultra-metric d which generates the topology of X, whereas the eigenfunctions depend on d and μ.

Let us now return to outlines of the two major applications of our theory, as indicated above in items (G) and (H).

First of all, we show that the p-adic Laplace operator introduced by Vladimirov and his collaborators fits into this setting as one of our possible Markov generators. As an important step towards this goal, we first relate our approach to the theory of p-adic fractional derivatives, as introduced by Vladimirov and collaborators [44]. This implies simple direct proofs of many results of [44], without using Fourier Analysis and the theory of Bruhat distributions.

The other main application has a slightly different flavour. We first outline a structure-theoretical feature, namely, the known fact that every locally compact ultra-metric space arises as the boundary of a locally finite *tree.* In particular, if the space is compact, then we can describe it as the boundary of a rooted tree.

Discrete time Random walks of nearest neighbour type on the vertex set of a tree are very well understood. See Chapter 9 of the (little read) book by WOESS [47].

It is then natural to ask if there is a connection between such random walks on trees and isotropic jump processes of the form (1.4) on the boundary of a tree.

Indeed, in recent work, KIGAMI [24] starts with a transient nearest neighbour random walk on a tree and constructs a naturally associated jump process on the boundary of the tree. Using this approach, he undertakes a detailed analysis of the process on the boundary. Restricting attention to the compact case, in the next section we shall answer the obvious question how the approaches of Kigami and of the present paper are related: the relation is basically one-to-one. As in [24], we take the ultra-metric space to be the geometric boundary at infinity of a locally finite rooted tree where each vertex has at least two forward neighbors.

"Basically" means that we need to restrict to random walks on trees which are *Dirichlet regular,* that is, the Dirichlet problem at infinity admits solution, or equivalently, the Green kernel of the random walk vanishes at infinity. (We comment on this condition, which appears to be natural in the present context, at the end.)

Given such a nearest neighbour random walk on a tree, its reversibility leads to an interpretation of the tree as an infinite electric network. This comes along with a natural Dirichlet form on the tree, and a natural approach is to use the Dirichlet form on the boundary which reproduces the power ("energy") of harmonic functions on the tree via their boundary values. This form on the boundary is computed with some effort in [24]; it induces the jump process studied there. Now, that form on the boundary is an integral with respect to the *Naïm kernel,* which goes back to the work of NAÏM [30] and DOOB [16] in the setting of abstract potential theory on spaces which are locally Euclidean. Trees do not have the latter property, but the validity of the resulting formula for the power of harmonic functions is proved for general infinite electric networks ($\equiv$ reversible random walks) in a forthcoming paper of KAIMANOVICH AND

Georgakopoulos [19]. A direct and rather simple proof for the case of trees is also given in §10.

We relate the Dirichlet form on the space of harmonic functions with finite power with the Dirichlet form on the boundary that is computed via the Naïm kernel in terms of the boundary values of the involved harmonic functions. This leads to an explanation of the relation between our isotropic jump processes of and the processes of [24] induced on the boundary of a tree by a random walk on that tree: every boundary process induced by a random walk is an isotropic jump process of our setting. Conversely, we show that up to a unique linear time change, every isotropic jump process on the boundary of a tree arises from a uniquely determined random walk as the process of [24]. Finally, in §12 we work out in a specific example how the applications of (G) and (H) are related: we consider the p-adic fractional derivative on the (compact) group of p-adic integers and the corresponding random walk on the associated tree.

2 Heat semigroup and heat kernel

We always consider a proper ultra-metric space (X, d) with the properties (i), (ii) outlined in the Introduction, and let μ be a Radon measure supported by the whole of X. We assume that

$$\begin{aligned} &\bullet\ \ \mu(\{x\}) = 0 \ \text{ if } x \in X \ \text{ is not an isolated point.}\\ &\bullet\ \ \mu(\{x\}) > 0 \ \text{ if } x \in X \ \text{ is an isolated point.} \end{aligned} \tag{2.1}$$

In most typical cases where (X, d) is non-compact, one will have $\mu(X) = \infty$.

We set

$$\Lambda(x) = \Lambda_d(x) = \{d(x, y) : y \in X\} \quad \text{and} \quad \Lambda = \Lambda_d = \bigcup_x \Lambda(x).$$

Each set $\Lambda(x)$ is countable. There is a strictly increasing sequence of numbers $r_k = r_k(x) > 0$ such that

$$\begin{aligned}
&\bullet \text{ if } (X,d) \text{ is non-compact and } x \text{ is not isolated then} && \Lambda(x) = && \{0\} \cup \{r_k : k \in \mathbb{Z}\}\\
&\qquad \text{with } \lim_{k\to\infty} r_k = \infty \text{ and } \lim_{k\to-\infty} r_k = 0; && && \\
&\bullet \text{ if } (X,d) \text{ is non-compact and } x \text{ is isolated then} && \Lambda(x) = && \{0\} \cup \{r_k : k \in \mathbb{Z}^+\}\\
&\qquad \text{with } \lim_{k\to\infty} r_k = \infty; && && \\
&\bullet \text{ if } (X,d) \text{ is compact and } x \text{ is not isolated then} && \Lambda(x) = && \{0\} \cup \{r_k : k \in \mathbb{Z}^-\}\\
&\qquad \text{with } \lim_{k\to-\infty} r_k = 0. && && \\
&\bullet \text{ if } (X,d) \text{ is compact and } x \text{ is isolated then} && \Lambda(x) = && \{0\} \cup \{r_1, \dots, r_n\}\\
&\qquad \text{is a finite set, with } n = n_x. && &&
\end{aligned} \tag{2.2}$$

The following notion will be used several times in later sections.

Definition 2.1 Given a closed ball $B = B_r(x) \subsetneq X$, its *predecessor* is the ball $B' = B_{r'}(x)$, where r' is the minimal element of $\Lambda(x)$ that satisfies $r' > r$.

Regarding the distance distribution σ on the half-line that appears in the definition (1.3) of P, we shall always denote its (left-continuous) distribution function by $\sigma(r) = \sigma\big([0\,,\,r)\big)$ and assume that it has the following non-degeneracy properties.

$$\sigma(0+) = 0\,, \quad \text{and} \quad \sigma(r) \ \text{ is strictly increasing and } < 1 \ \text{ on each } \ \Lambda(x)\,,\ x \in X. \tag{2.3}$$

The assumptions (2.3) imply that $\sigma(r) > 0$ for every $r > 0$ and that σ charges every interval $[r\,,\,r + \varepsilon)$, where $\varepsilon > 0$ and $r \in \Lambda$.

Recall that σ^t denotes the probability distribution with distribution function $\sigma(r)^t$. Also recall the operator Q_r from (1.2),

$$\mathrm{Q}_r f(x) = \frac{1}{\mu(B_r(x))} \int_{B_r(x)} f \, d\mu.$$

Since

$$B_r(x) = B_r(y) \quad \text{for every } y \in B_r(x), \tag{2.4}$$

we have that $\mathrm{Q}_r f$ takes constant values on any ball $B_r(x)$ with given radius $r > 0$. In particular, $x \mapsto \mathrm{Q}_r f(x)$ is a bounded continuous function. We can write $\mathrm{Q}_r f$ in the form

$$\mathrm{Q}_r f(x) = \int_X K_r(x,y) f(y) d\mu(y),$$

where the integral kernel is given by the equation

$$K_r(x,y) = \frac{1}{\mu(B_r(x))} \mathbf{1}_{B_r(x)}(y). \tag{2.5}$$

By (2.4), the kernel is symmetric:

$$K_r(x,y) = K_r(y,x).$$

Since $\mathrm{Q}_r f \geq 0$ when $f \geq 0$, and $\mathrm{Q}_r \mathbf{1} = \mathbf{1}$, the operator Q_r is symmetric and Markovian. It follows that $\mathrm{Q}_{B_r} f$ extends as a bounded self-adjoint operator acting on $L^2(X,\mu)$ with norm $\|\mathrm{Q}_r\|_{L^2 \to L^2} = 1$. Moreover, for any $r, r' > 0$,

$$\mathrm{Q}_{r'} \mathrm{Q}_r = \mathrm{Q}_r \mathrm{Q}_{r'} = \mathrm{Q}_{\max\{r,r'\}} \,. \tag{2.6}$$

Thus $\{\mathrm{Q}_r\}_{r \geq 0}$ is a non-increasing family of ortho-projectors. We have $\mathrm{Q}_0 = id$. By monotonicity, the strong limit $\mathrm{Q}_\infty = \lim_{r \to \infty} \mathrm{Q}_r$ exists on L^2 and is again an ortho-projector. If (X,d) is non-compact, it is easy to see that $\mathrm{Q}_\infty = 0$. If (X,d) is compact then Q_∞ coincides with the ortho-projector on the one dimensional subspace of L^2 consisting of constant functions. Consequently, the family of ortho-projectors $\{E_\tau\}_{-\infty}^{+\infty}$, defined as

$$E_\tau = \begin{cases} \mathrm{Q}_{B_{1/\tau}} & \text{if } \ \tau \geq 0 \\ 0 & \text{if } \ \tau < 0 \end{cases}$$

defines a spectral measure dE_τ in L^2. What is crucial in our analysis is that the measure dE_τ gives the spectral resolution of the Markov semigroup (P^t) defined in the Introduction in (1.4). We now verify its basic properties.

Theorem 2.2 (1) *The family* $\{P^t\}_{t \geq 0}$ *defined in (1.4) is a symmetric Markov semigroup on* $L^2(X,\mu)$.

(2) *Under the assumptions (2.3) on* σ*, the semigroup is strongly continuous.*

Proof. For any given $s, t > 0$ and $f \in L^2$, we have by (2.6)

$$\begin{aligned} P^s P^t f(x) &= \int_0^\infty d\sigma^s(r) \int_0^\infty d\sigma^t(r') \, \mathrm{Q}_r \mathrm{Q}_{r'} f(x) = \\ &= \int_0^\infty d\sigma^s(r) \int_0^\infty d\sigma^t(r') \, \mathrm{Q}_{\max\{r,r'\}} f(x). \end{aligned}$$

Let ξ_1 and ξ_2 be two independent random variables with distributions σ^s and σ^t, respectively. Then the distribution of the random variable $\xi = \max\{\xi_1, \xi_2\}$ is σ^{t+s}. It follows that

$$P^s P^t f(x) = \mathbb{E}\Big(\mathrm{Q}_{\max\{\xi_1,\xi_2\}} f(x)\Big) = \int_0^\infty \mathrm{Q}_r f(x)\, d\sigma^{t+s}(r) = P^{t+s} f(x).$$

This proves the first statement.

For the second statement, we already observed that $\mathrm{Q}_r f \to f$ strongly as $r \to 0$. Fix $\varepsilon > 0$ and choose $\delta_1 > 0$ such that

$$\sup_{0 \le r < \delta_1} \|\mathrm{Q}_r f - f\|_{L^2} < \frac{\varepsilon}{2}.$$

Since $\sigma(r) > 0$ for all $r > 0$, we can choose $\delta_2 > 0$ such that

$$\sup_{0 < t < \delta_2} \|f\|_{L^2} \Big((1 - \sigma^t(\delta_2)\Big) < \frac{\varepsilon}{4}.$$

Thus, for all $0 < t < \delta_2$ we obtain

$$\begin{aligned} \left\|\mathrm{Q}^t f - f\right\|_{L^2} &\le \int_{[0,\delta_1)} \|\mathrm{Q}_r f - f\|_{L^2}\, d\sigma^t(r) + 2\,\|f\|_{L^2}\,(1 - \sigma^t(\delta_2)) \le \\ &\le \sup_{0 \le r < \delta_1} \|\mathrm{Q}_r f - f\|_{L^2} + \frac{\varepsilon}{2} < \varepsilon, \end{aligned}$$

whence the semigroup $\{P^t\}$ is strongly continuous. ■

Remark 2.3 We cannot relax all the assumptions (2.3) on σ. In particular, when X is non-discrete, then it is necessary for strong continuity of $\{P^t\}$ that $\sigma(r) > 0$ for all $r > 0$. Indeed, if we assume that $\sigma(r) = 0$ on some interval $[0\,,\,b]$, then for any $s < b$,

$$\begin{aligned} \mathrm{Q}_s P^t f &= \int_{[b\,,\,\infty)} \mathrm{Q}_r \mathrm{Q}_s f\, d\sigma^t(r) \\ &= \int_{[b\,,\,\infty)} \mathrm{Q}_{\max\{r,s\}} f\, d\sigma(r)^t = P^t f. \end{aligned}$$

This equality would contradict the continuity of the semigroup $\{P^t\}$. Indeed letting $t \to 0$ and assuming that $P^t f \to f$ we would come to the conclusion that $\mathrm{Q}_s f = f$ for any function $f \in L^2$ and any $s < b$. This is not true when X is non-discrete, since in that case, each distance set $\Lambda(x)$ contains arbitrarily small positive numbers.

We shall from now on always stick to our assumptions (2.1) and (2.3) on μ and σ.

Theorem 2.4 *The operator* P^t *admits an integral kernel* $p(t, x, y)$, *that is,*

$$P^t f(x) = \int_X p(t, x, y) f(y) d\mu(y).$$

We call it the heat kernel. *It is given explicitly as*

$$p(t, x, y) = \int_{\{r \ge d(x,y)\}} \frac{d\sigma^t(r)}{\mu\big(B_r(x)\big)}. \tag{2.7}$$

Proof. By the assumption that $\sigma(0+) = 0$, and using Fubini's theorem,

$$\begin{aligned}
P^t f(x) &= \int_{\{r\geq 0\}} \mathrm{Q}_r f(x)\, d\sigma^t(r) \\
&= \int_{\{r>0\}} \left(\frac{1}{\mu(B_r(x))} \int_X \mathbf{1}_{B_r(x)} f(y)\, d\mu(y) \right) d\sigma^t(r) \\
&= \int_X \left(\int_{\{r\geq d(x,y)\}} \frac{1}{\mu(B_r(x))} f(y) \right) d\sigma^t(r)\, d\mu(y)\,,
\end{aligned}$$

which concludes our computation. ■

3 Spectral Distribution Function

We continue to work with the (d, μ, σ)-process and the associated symmetric Markov semigroup $\{P^t\}_{t>0}$ on our proper ultra-metric space X, always assuming that μ and σ satisfy (2.1) and (2.3). Recall the intrinsic ultra-metric defined in (1.6), that is, $d_*(x, y) = -1/\log \sigma(r)$ whenever $d(x, y) = r$. Since $\sigma(r) < 1$ by (2.3), this is well-defined.

Clearly, the value sets of the new metric are

$$\Lambda_*(x) = \{d_*(x, y) : y \in X\} = \{-1/\log \sigma(r) : r \in \Lambda(x)\}\,, \quad \text{and} \quad \Lambda_* = \bigcup_x \Lambda_*(x)\,. \tag{3.1}$$

The following is straightforward.

Lemma 3.1 *For $r \in \Lambda(x)$, resp. $r^* \in \Lambda_*(x)$, we have*

$$B^*_{r^*}(x) = B_r(x) \Longleftrightarrow \frac{1}{r^*} = \log \frac{1}{\sigma(r)}\,,$$

where $B^*_{r^*}(x)$ is the closed d_*-ball of radius r^* centred at x.

In particular, the families of closed d-balls and of closed d_*-balls coincide.

Definition 3.2 For any $x \in X$ we define the *spectral distribution function* $N : [0\,,\,\infty) \to [0\,,\,\infty)$ as

$$N(x, \tau) = 1/\mu\big(B^*_{1/\tau}(x)\big). \tag{3.2}$$

See Figures 1, 2 and 3 for qualitative pictures.

Theorem 3.3 *For $x, y \in X$,*

$$p(t, x, y) = t \int_0^{1/d_*(x,y)} N(x, \tau) \exp(-\tau t)\, d\tau. \tag{3.3}$$

In particular,

$$p(t, x, x) = \int_{[0,\infty)} \exp(-\tau t)\, dN(x, \tau) \tag{3.4}$$

(Lebesgue integral with respect to the measure whose distribution function is $N(x, \cdot)$)



Figure 1: The graph of the function $\tau \mapsto N(x,\tau)$ in the case when (X,d) is compact



Figure 2: The graph of the function $\tau \mapsto N(x,\tau)$ in the case, when (X,d) is discrete and infinite



Figure 3: The graph of the function $\tau \mapsto N(x,\tau)$ in the case when (X,d) is not discrete and non-compact

Proof. Let $\Lambda(x) = \{r_k\}$ as in (2.2), where k ranges in $\mathbb{Z}$, $\mathbb{Z}^+$ or $\mathbb{Z}^-$, or the range of k is finite according to the respective case. Observe that on each interval $[r_k\,,\,r_{k+1})$, we have $B_r(x) = B_{r_k}(x)$. Thus, the function $x \mapsto \mu\big(B_r(x)\big)$ takes constant values on that interval.

Let y be such that $d(x,y) = r_n$. By Theorem 2.7 we can write

$$\begin{aligned} p(t,x,y) &= \int_{\{r\geq r_n\}} \frac{d\sigma^t(r)}{\mu\big(B_r(x)\big)} = \sum_{k\geq n} \int_{[r_k,r_{k+1})} \frac{d\sigma^t(r)}{\mu\big(B_r(x)\big)} \\ &= \sum_{k\geq n} \frac{\sigma(r_{k+1})^t - \sigma(r_k)^t}{\mu\big(B_{r_k}(x)\big)} \\ &= \sum_{k\geq n} \frac{1}{\mu\big(B_{r_k}(x)\big)} \int_{\sigma(r_k)}^{\sigma(r_{k+1})} t\xi^{t-1}\,d\xi. \end{aligned}$$

Now let

$$\tau(r) = \log\frac{1}{\sigma(r)}\,, \quad 0 \leq r \leq \infty.$$

By the assumptions (2.3) on σ,

$$\tau(r_{k+1}) < \tau(r_k) \quad \text{and} \quad \tau(r_{k+1}) \leq \tau(r) \leq \tau(r_k) \quad \text{for } \ r_k \leq r \leq r_{k+1}\,.$$

By Lemma 3.1, with the substitution $\xi = e^{-\tau}$, and using that $B^*_{1/\tau(r_k)}(x) = B^*_{1/\tau}(x)$ for every τ with $\tau(r_{k+1}) < \tau < \tau(r_k)$,

$$\begin{aligned} \frac{1}{\mu\big(B_{r_k}(x)\big)} \int_{\sigma(r_k)}^{\sigma(r_{k+1})} t\xi^{t-1}\,d\xi &= \frac{1}{\mu\big(B^*_{1/\tau(r_k)}(x)\big)} \int_{\tau(r_{k+1})}^{\tau(r_k)} t\exp(-\tau t)\,d\tau \\ &= \int_{\tau(r_{k+1})}^{\tau(r_k)} \frac{t\exp(-\tau t)}{\mu\big(B^*_{1/\tau}(x)\big)}\,d\tau = t\int_{\tau(r_{k+1})}^{\tau(r_k)} N(x,\tau)\,\exp(-\tau t)\,d\tau. \end{aligned}$$

Summing over all $k \geq n$, the identity (3.3) follows. The second identity follows from the first one by integration by parts. ■

Let us define $t_* = t_*(x) \leq \infty$ as follows.

$$t_* = \limsup_{\tau\to\infty} \frac{\log N(x,\tau)}{\tau}\,. \tag{3.5}$$

It controls the exponential decay of the intrinsic volume function $V(x,r)$ of (1.7), as $r \to 0$.

Corollary 3.4 (a) *For any x, y and $t > 0$,*

$$0 < p(t,x,y) < \min\{p(t,x,x), p(t,y,y)\} \leq \infty\,.$$

(b) $p(t,x,x) = \infty$ *for* $0 < t < t_*\,,$ *and* $p(t,x,x) < \infty$ *for* $t > t_*\,.$

Proposition 3.5 *Assume that (X,d) is compact and that $t_*(x) < \infty$ for some $x \in X$. Then uniformly in $y \in X$,*

$$\lim_{t\to\infty} p(t,x,y) = \frac{1}{\mu(X)}.$$

Proof. We have $\tau_0 > 0$, where $1/\tau_0$ is the d_*-diameter of X. Hence $N(x,\tau) = 1/\mu(X)$ for $0 < \tau < \tau_0$. We write

$$\begin{aligned} p(t,x,x) &= t\int_0^\infty N(x,\tau)\exp(-\tau t)\,d\tau = \\ &= \frac{1}{\mu(X)}\big(1 - \exp(-\tau_0 t)\big) + t\int_{\tau_0}^\infty N(x,\tau)\exp(-\tau t)\,d\tau. \end{aligned}$$

Since $t_*(x) < \infty$, we can choose $A, a > 0$ such that $N(x,\tau) \le A\exp(a\tau)$ for all $\tau > \tau_0$. Then for all $t > 2a$ we obtain

$$\begin{aligned} t\int_{\tau_0}^{\infty} N(x,\tau)\exp(-\tau t)\,d\tau &\le \frac{At}{t-a}\exp(-\tau_0(t-a)) \\ &\le 2A\exp\left(-\frac{\tau_0\, t}{2}\right). \end{aligned}$$

It follows that

$$\lim_{t\to\infty} p(t,x,x) = \frac{1}{\mu(X)}. \tag{3.6}$$

For arbitrary $y \in X$ we know that

$$p(t,x,y) \le p(t,x,x), \tag{3.7}$$

and

$$\begin{aligned} p(t,x,y) &\ge p(t,x,x) - 2\exp\left(-\frac{t}{2d_*(x,y)}\right) p\left(\frac{t}{2},x,x\right) \ge \\ &\ge p(t,x,x)\left(1 - 2\exp\left(-\frac{\tau_0\, t}{2}\right)\frac{p(\frac{t}{2},x,x)}{p(t,x,x)}\right). \end{aligned} \tag{3.8}$$

Equations (3.6), (3.7) and (3.8) yield the result. ■

Proposition 3.6 *Assume that* (X,d) *is non-compact and that* $t_*(x) < \infty$ *for some* $x \in X$. *Then*

$$\lim_{t\to\infty}\frac{p(t,x,y)}{p(t,x,x)} = 1,$$

locally uniformly in y.

Proof. Choose $A, a > 0$ such that $N(x,\tau) \le A\exp(a\tau)$ for all $\tau > 0$. For $t > 2a$ and $y \in X$ we write

$$\begin{aligned} p(t,x,x) - p(t,x,y) &= t\int_{1/d_*(x,y)}^{\infty} N(x,\tau)\,\exp(-\tau t)\,d\tau \\ &\le At\int_{1/d_*(x,y)}^{\infty} \exp\bigl(-\tau(t-a)\bigr)\,d\tau \le 2A\exp\left(-\frac{t-a}{d_*(x,y)}\right). \end{aligned} \tag{3.9}$$

On the other hand

$$\begin{aligned} p(t,x,x) &= \int_{[0\,,\infty)} \exp(-\tau t)\,dN(x,\tau) \ge \\ &\ge \int_{[0\,,\,1/2d_*(x,y))} \exp(-\tau t)\,dN(x,\tau) \ge \exp\left(-\frac{t}{2d_{*(x,y)}}\right) N\left(x,\frac{1}{2d_*(x,y)}\right). \end{aligned} \tag{3.10}$$

Let K be a compact set and $D(K)$ be its d_*-diameter. The inequalities 3.9 and 3.10 yield the following: for $y \in K$,

$$\begin{aligned} \frac{p(t,x,x) - p(t,x,y)}{p(t,x,x)} &\le \frac{2A\exp\left(-\frac{t-a}{d_*(x,y)}\right)}{\exp\left(-\frac{t}{2d_*(x,y)}\right) N\left(x,\frac{1}{2d_*(x,y)}\right)} \\ &\le \frac{2A\exp\left(-\frac{t-2a}{2D(K)}\right)}{N\left(x,\frac{1}{2D(K)}\right)} \to 0 \text{ as } t\to\infty. \end{aligned}$$

This completes the proof. ■

4 Basic estimates of the Heat Kernel

The purpose of this section is to provide estimates of the heat kernel (transition density)

$$p(t,x,y) = t\int_0^{1/d_*(x,y)} N(x,\tau)\exp(-\tau t)\,d\tau$$

of (3.3) in Theorem 3.3.

For any non-decreasing real function N set

$$\widehat{N}(t,r) = t\int_0^{r} N(\tau)\exp(-t\tau)\,d\tau. \tag{4.1}$$

Then for any fixed x, the heat kernel $p(t,x,y)$ can be written in the form

$$p(t,x,y) = \widehat{N}\left(t, \frac{1}{d_*(x,y)}\right), \tag{4.2}$$

where $N(\tau) = N(x,\tau)$ is the spectral distribution function at x, as introduced in Definition 3.2. Our heat kernel estimates will be based on studying the function $\widehat{N}(t,r)$ as a functional of N.

Proposition 4.1 *For all $x, y \in X$, the following estimates hold.*

(a) *For all $0 < t \le d_*(x,y)$,*

$$\frac{t}{2e\,d_*(x,y)} N\left(x, \frac{1}{2d_*}\right) \le p(t,x,y) \le \frac{t}{d_{*(x,y)}} N\left(x, \frac{1}{d_*}\right). \tag{4.3}$$

(b) *For all $t > d_*(x,y)$,*

$$\frac{1}{2e} N\left(x, \frac{1}{2t}\right) \le p(t,x,y).$$

(c) *For all $t > 0$,*

$$\frac{1}{e} N\left(x, \frac{1}{t}\right) \le p(t,x,x).$$

Proof. For (a), we use monotonicity of $\tau \mapsto N(x,\tau)$ and write

$$\begin{aligned}
p(t,x,y) &= t\int_0^{1/d_*(x,y)} N(x,\tau)\exp(-\tau t)\,d\tau \le \frac{t}{d_*(x,y)} N\left(x, \frac{1}{d_*(x,y)}\right) \qquad \text{and} \\
p(t,x,y) &\ge t\int_{1/2d_*(x,y)}^{1/d_*(x,y)} N(x,\tau)\exp(-\tau t)\,d\tau \\
&\ge \frac{t}{2d_*(x,y)} N\left(x, \frac{1}{2d_*(x,y)}\right)\exp(-\frac{t}{d_*(x,y)}) \ge \frac{t}{2e\,d_*(x,y)} N\left(x, \frac{1}{2d_*(x,y)}\right).
\end{aligned}$$

For (b), when $1/d_*(x,y) > 1/t$ we obtain

$$\begin{aligned}
p(t,x,y) &\ge t\int_0^{1/t} N(x,\tau)\exp(-\tau t)\,d\tau \\
&\ge t\int_{1/2t}^{1/t} N(x,\tau)\exp(-\tau t)\,d\tau \ge \frac{1}{2e} N\left(x, \frac{1}{2t}\right).
\end{aligned}$$

For (c), using (3.4) we write

$$\begin{aligned} p(t,x,x) &= t\int_0^\infty N(x,\tau)\exp(-\tau t)\,d\tau \geq t\int_{1/t}^\infty N(x,\tau)\exp(-\tau t)\,d\tau \\ &\geq N\left(x,\frac{1}{t}\right)\int_1^\infty \exp(-\tau)d\tau = \frac{1}{e}N\left(x,\frac{1}{t}\right). \end{aligned}$$

The proof is finished. ■

Definition 4.2 A non-decreasing function $\Phi : \mathbb{R}_+ \to \mathbb{R}_+$ is said to satisfy the *doubling property* if there exists a constant $D > 0$ such that

$$\Phi(2s) \leq D \cdot \Phi(s) \quad \text{for all } s > 0.$$

It is known (Potter's theorem) that if Φ is doubling then

$$\Phi(s_2) \leq D\left(\frac{s_2}{s_1}\right)^\delta \Phi(s_1) \quad \text{for all } 0 < s_1 < s_2\,, \text{ where } \delta = \log_2 D. \tag{4.4}$$

Proposition 4.3 *For any given $x \in X$, the following two properties are equivalent.*

(1) For some constant $c > 0$ and all $t > 0$,

$$\frac{1}{c}N\left(x,\frac{1}{t}\right) \leq p(t,x,x) \leq c\,N\left(x,\frac{1}{t}\right). \tag{4.5}$$

(2) The function $\tau \mapsto N(x,\tau)$ is doubling.

Proof. (2) $\Rightarrow$ (1): The lower bound is the inequality (4.1), so what is left is to prove the upper bound. We have

$$\begin{aligned} p(t,x,x) &= t\int_0^\infty N(x,\tau)\exp(-\tau t)\,d\tau \\ &= \int_0^\infty N\left(x,\frac{\tau}{t}\right)\exp(-\tau)\,d\tau. \end{aligned}$$

By assumption, N is doubling, whence

$$\begin{aligned} \int_0^\infty N\left(x,\frac{\tau}{t}\right)\exp(-\tau)\,d\tau &= N\left(x,\frac{1}{t}\right)\int_0^\infty \frac{N\left(x,\frac{\tau}{t}\right)}{N\left(x,\frac{1}{t}\right)}\exp(-\tau)\,d\tau \\ &\leq D\,N\left(x,\frac{1}{t}\right)\int_0^\infty \max\{1,\tau^\delta\}\exp(-\tau)\,d\tau \\ &\leq D'\,N\left(x,\frac{1}{t}\right). \end{aligned}$$

The constants $D, D', \delta > 0$ come from Potter's bound (4.4).

(1) $\Rightarrow$ (2): Using the inequality (4.5) we obtain

$$\begin{aligned} c\,N\left(x,\frac{1}{t}\right) &\geq p(t,x,x) \geq t\int_{2/t}^\infty N(x,\tau)\exp(-\tau t)\,d\tau \\ &\geq e^{-2}N(x,2/t). \end{aligned}$$

Setting $\tau = 1/t$, we come to the desired conclusion. ■

Theorem 4.4 *The following inequalities hold for all $x, y \in X$ and $t > 0$.*

$$p(t,x,y) \geq \frac{1}{2e}\,\frac{t}{t+d_{*(x,y)}}\,N\left(x, \frac{1}{2\big(t+d_*(x,y)\big)}\right), \tag{4.6}$$

$$p(t,x,y) \leq 2e\,\frac{t}{t+d_*(x,y)}\,N\left(\frac{t+d_*(x,y)}{2}, x, x\right), \tag{4.7}$$

Proof. When $0 < t \leq d_*(x,y)$ we use Proposition 4.1(a):

$$\begin{aligned} p(t,x,y) &\geq \frac{1}{2e}\frac{t}{d_*(x,y)}N\left(x, \frac{1}{2d_*(x,y)}\right) \\ &\geq \frac{1}{2e}\frac{t}{t+d_*(x,y)}N\left(x, \frac{1}{2\big(t+d_*(x,y)\big)}\right). \end{aligned}$$

When $t > d_*$ we use Proposition 4.1(b):

$$\begin{aligned} p(t,x,y) &\geq \frac{1}{2e}N\left(x, \frac{1}{2t}\right) \\ &\geq \frac{1}{2e}\frac{t}{t+d_*(x,y)}N\left(x, \frac{1}{2\big(t+d_*(x,y)\big)}\right). \end{aligned}$$

Similarly for the upper bound (4.7): depending on whether $0 < t \leq d_*(x,y)$, or $t > d_*(x,y)$, we will have

$$\begin{aligned} \frac{t}{t+d_*(x,y)}\,p\left(\frac{t+d_*(x,y)}{2}, x, x\right) &\geq \frac{t}{2d_*(x,y)}\,p\big(d_*(x,y), x, x\big) \\ &\geq \frac{1}{2e}\,\frac{t}{d_*(x,y)}N\left(x, \frac{1}{d_*(x,y)}\right) \geq \frac{1}{2e}\,p(t,x,y), \end{aligned}$$

or

$$\frac{t}{t+d_*(x,y)}\,p\left(\frac{t+d_*(x,y)}{2}, x, x\right) \geq \frac{1}{2e}\,p(t,x,y),$$

respectively, as desired. ■

Corollary 4.5 *Suppose that $\tau \mapsto N(x,\tau)$ is doubling. Then*

$$p(t,x,y) \leq \frac{C\,t}{t+d_*}\,N\left(x, \frac{1}{t+d_*}\right), \tag{4.8}$$

$$p(t,x,y) \geq \frac{c\,t}{t+d_*}\,N\left(x, \frac{1}{t+d_*}\right), \tag{4.9}$$

for all $t > 0, y \in X$ and some $C, c > 0$.

The following three cases are of particular interest.

Example 4.6 Assume that (X,d) is non-compact and has no isolated points, e.g. $X = \mathbb{Q}_p$. Assume that for a given $x \in X$, and some $\alpha, \beta > 0$,

$$N(x,\tau) \simeq \begin{cases} \tau^{\alpha}, & 0 < \tau \leq 1, \\ \tau^{\beta}, & \tau > 1. \end{cases}$$

Then for all $t > 0$ and all $y \in X$,

$$p(t,x,y) \simeq \begin{cases} t\big(t+d_*(x,y)\big)^{-1-\beta}, & 0 < t+d_*(x,y) \leq 1, \\ t\big(t+d_*(x,y)\big)^{-1-\alpha}, & t+d_*(x,y) > 1. \end{cases}$$

Example 4.7 Assume that (X, d) is discrete, e.g. $X = \bigoplus_{k=1}^{\infty} \mathbb{Z}(n_k)$. Assume that for a given $x \in X$,

$$N(x, \tau) \simeq \tau^{\gamma}$$

for some $\gamma > 0$ and all $0 < \tau \leq 1$. Then for all $t > 0$ and all $y \in X$ such that $t + d_* \geq 1$,

$$p(t, x, y) \simeq \frac{t}{\big(t + d_*(x, y)\big)^{1+\gamma}}.$$

Example 4.8 Assume that (X, d) is compact and has no isolated points, e.g. $X = \mathbb{Z}_p$, the set of p-adic integers. Assume that for a given $x \in X$,

$$N(x, \tau) \simeq \tau^{\delta},$$

for some $\delta > 0$ and all $\tau > 1$. Then for all $t > 0$ and all $y \in X$ such that $0 < t + d_*(x, y) \leq 1$,

$$p(t, x, y) \simeq \frac{t}{\big(t + d_*(x, y)\big)^{1+\delta}}.$$

Thus in all three examples $p(t, x, y)$ has a shape similar to that of the Cauchy density in Euclidean space. The parameters α, β, γ and δ play the role of a local dimension (resp. dimension at infinity) of the space X.

Definition 4.9 A non-decreasing function $\Psi : \mathbb{R}_+ \to \mathbb{R}_+$ is said to satisfy the *reverse doubling property,* if

$$\Psi(r) \geq (1 + \eta)\Psi(\delta r),$$

for all $r > 0$ and some $\delta, \eta \in (0\,,\,1)$.

The function is said to satisfy the reverse doubling condition at 0, resp. at ∞, if there is $R > 0$ such that the above inequality holds for all r with $0 < r < R$, resp. all $r > R$.

Note that Ψ satisfies the reverse doubling property if and only if its generalized inverse Ψ^{-1} is doubling, where $\Psi^{-1}(s) = \sup\{r : \Psi(r) \leq s\}$. We have the following sufficient criterion for $N(x, \cdot)$ to have the doubling property.

Proposition 4.10 *The function $\tau \mapsto N(x, \tau)$ is doubling if the following two conditions hold:*

1. *The function $\Psi(r) = -1/\log \sigma(r)$ satisfies the reverse doubling property.*
2. *The volume function $r \mapsto V(x, r) = \mu\big(B_r(x)\big)$ satisfies the doubling property.*

Proof. It follows from the Definition 3.2 of the spectral distribution function that $\tau \mapsto N(x, \tau)$ is doubling if and only if the function $s \mapsto \mu(B^*_s(x))$ is doubling.

By definition, resp. Lemma 3.1,

$$B^*_s(x) = B_r(x),\,,\ r = \Psi^{-1}(s)\,, \quad \text{and} \quad B^*_{2s}(x) = B_{r'}(x)\,,\ r'^{-1}(2s).$$

Condition (2) implies that for some constants $A, a > 0$,

$$\mu\big(B^*_{2s}(x)\big) = \mu\big(B_{r'}(x)\big) \leq A\,(r'/r)^a\,\mu\big(B_r(x)\big) = A\,(r'/r)^a\,\mu\big(B^*_s(x)\big).$$

To estimate the ratio $(r'/r)^a$ we use condition (1):

$$\frac{1}{2}\log\frac{1}{\sigma(r)} = \log\frac{1}{\sigma(r')} \leq \frac{1}{1+\eta}\log\frac{1}{\sigma(\delta r')} \leq \cdots \leq \frac{1}{(1+\eta)^k}\log\frac{1}{\sigma(\delta^k r')}.$$

We choose $k > 1$ such that $(1+\eta)^{-k} < 1/2$. For such k we obtain

$$\log \frac{1}{\sigma(r)} < \log \frac{1}{\sigma(\delta^k r')}.$$

It follows that $r \geq \delta^k r'$, whence

$$\mu\big((B^*_{2s}(x)\big) \leq A\,\delta^{-ak}\mu\big(B^*_s(x)\big).$$

The proof is finished. ∎

Example 4.11 Let (X, d) be the field of p-adic numbers $\mathbb{Q}_p$ equipped with its canonical ultrametric $d(x,y) = \|x-y\|_p$. Let μ be the additive Haar measure on $\mathbb{Q}_p$. Let $\mathbb{Z}_p \subset \mathbb{Q}_p$ be the group of p-adic integers. For $x \in \mathbb{Q}_p$, $k \in \mathbb{Z}$ and $p^k \leq r < p^{k+1}$, we have

$$B_r(x) = x + p^{-k}\mathbb{Z}_p \quad \text{and} \quad \mu\big(B_r(x)\big) = p^k. \tag{4.10}$$

In particular, for all $r > 0$,

$$\frac{r}{p} < \mu\big(B_r(x)\big) \leq r.$$

Let us choose the distance distribution with distribution function $\sigma(r) = \exp\big(-(p/r)^\alpha\big)$, where $\alpha > 0$. Then $\Psi(r) = 1/\log \frac{1}{\sigma(r)} = (r/p)^\alpha$. By Proposition 4.10, the function $\tau \to N(x,\tau)$ is doubling. Direct computations show that

$$d_*(x,y) = \left(\frac{\|x-y\|_p}{p}\right)^\alpha \quad \text{and} \quad \lambda^{\frac{1}{\alpha}} < \mu\big(B^*_\lambda(x)\big) \leq p\, r^{1/\alpha},$$

whence

$$\frac{1}{p}\tau^{1/\alpha} \leq N(x,\tau) < \tau^{1/\alpha}.$$

By Corollary 4.5, there exist constants $c_1, c_2 > 0$ such that for all $t > 0$ and all $x, y \in \mathbb{Q}_p$,

$$\frac{c_1\, t}{(t^{1/\alpha} + \|x-y\|_p)^{1+\alpha}} \leq p(t,x,y) \leq \frac{c_2\, t}{(t^{1/\alpha} + \|x-y\|_p)^{1+\alpha}}.$$

On the other hand, choosing $\sigma(r) = \exp\Big(-\big(\log(1+\frac{1}{r})\big)^\beta\Big)$, where $\beta > 0$, we obtain

$$\frac{1}{p}\left(\exp(\tau^{1/\beta}) - 1\right) \leq N(x,\tau) \leq p\left(\exp(\tau^{1/\beta}) - 1\right).$$

Thus, the spectral distribution function $\tau \to N(x,\tau)$ is not doubling on the whole range of $\tau \in (0,\infty)$, and Corollary 4.5 does not apply. Note that $N(x,\tau) \simeq \tau^{1/\beta}$ at 0, whence we can apply Theorem 4.4 and the standard Laplace transform argument to show that

$$p(t,x,y) \simeq \frac{t}{(t^{1/\beta} + \|x-y\|_p)^{1+\beta}} \quad \text{when} \quad t + \|x-y\|_p \geq 1,$$

compare with Proposition 3.6. By other appropriate choices of the distance distribution σ, one can consider many other interesting examples.

5 Green function and transience

The resolvent $\{R_\lambda\}_{\lambda>0}$ associated with our (or any) semigroup $\{P^t\}_{t>0}$ is defined on the set of bounded Borel measurable functions f as

$$R_\lambda f = \int_0^\infty e^{-\lambda t} P^t f \, dt.$$

Since the semigroup admits an integral kernel $p(t,x,y)$ with respect to the reference measure μ, the resolvent admits an integral kernel $G_\lambda(x,y)$ as well. We call it the λ-*Green function.* It can be represented in the form

$$G_\lambda(x,y) = \int_0^\infty e^{-\lambda t} p(t,x,y)\, dt\,.$$

Theorem 5.1 *Let $\{P^t\}$ be defined by (1.4). Then its λ-Green function is given by*

$$G_\lambda(x,y) = \int_0^{1/d_*(x,y)} \frac{N(x,\tau)\, d\tau}{(\tau+\lambda)^2}\,. \tag{5.1}$$

The following properties hold:

(a) *The function $d_{**} : X \times X \to \mathbb{R}_+$ defined as*

$$d_{**}(x,y) = \begin{cases} 1/G_\lambda(x,y) & \text{if } \; x \neq y \\ 0 & \text{if } \; x = y \end{cases}$$

is an ultra-metric on X generating its topology.

(b) *For any given $\lambda > 0$,*

$$G_\lambda(x,y) \sim \lambda^{-2} \int_0^{1/d_*(x,y)} N(x,\tau)\, d\tau \quad \text{as } \; d_*(x,y) \to \infty\,.$$

(c) *Assume that $G_\lambda(x,x) = \infty$. Then*

$$G_\lambda(x,y) \sim \int_1^{1/d_*(x,y)} \frac{N(x,\tau)\, d\tau}{\tau^2} \quad \text{as } \; d_*(x,y) \to 0\,.$$

Proof. Using the definitions of R_λ and $G_\lambda(x,y)$, we obtain

$$R_\lambda f(x) = \int_0^\infty \int_X e^{-\lambda t} p(t,x,y) f(y)\, d\mu(y)\, dt = \int_X G_\lambda(x,y) f(y)\, d\mu(y),$$

so that $G_\lambda(x,y)$ is indeed the integral kernel of the operator R_λ. Integrating the identity (3.3) with respect to t, we obtain (5.1).

Let $d_*(x,z) = a$ and $d_*(y,z) = b$, assume that $a \leq b$. Then by the ultra-metric inequality, $d_*(x,y) \leq b$. Using the symmetry of the λ-Green function, we write

$$\begin{aligned} G_\lambda(x,z) &= \int_0^{1/a} \frac{N(z,\tau)\, d\tau}{(\tau+\lambda)^2} \geq \int_0^{1/b} \frac{N(z,\tau)\, d\tau}{(\tau+\lambda)^2} = G_\lambda(y,z) \qquad \text{and} \\ G_\lambda(x,y) &= \int_0^{1/d_*(x,y)} \frac{N(y,\tau)\, d\tau}{(\tau+\lambda)^2} \geq \int_0^{1/b} \frac{N(y,\tau)\, d\tau}{(\tau+\lambda)^2} = G_\lambda(y,z). \end{aligned}$$

It follows that

$$G_\lambda(x,y) \ge G_\lambda(y,z) = \min\{G_\lambda(x,z), G_\lambda(z,y)\},$$

whence the function $d_{**}(x,y) = 1/G_\lambda(x,y)$ satisfies the ultra-metric inequality. For any fixed $x \in X$, the new distance $d_{**}(x,y)$ is a strictly increasing function of $d_*(x,y)$. Therefore it generates the same topology as d_*. This proves (a).

Let $\lambda > 0$ be fixed and $d_*(x,y) \to \infty$. Using (5.1), we get

$$\begin{aligned} G_\lambda(x,y) &\ge \frac{1}{\left(1/d_*(x,y)+\lambda\right)^2} \int_0^{1/d_*(x,y)} N(x,\tau)\,d\tau \quad \text{and} \\ G_\lambda(x,y) &\le \frac{1}{\lambda^2} \int_0^{1/d_*(x,y)} N(x,\tau)\,d\tau. \end{aligned}$$

These two inequalities imply the asymptotic behavior proposed in (b). The proof of (c) is analogous. ■

Corollary 5.2 *Assume that the function $\tau \mapsto N(x,\tau)$ is doubling. Then we have the following.*

(1) *If $G_\lambda(x,x) = \infty$ then*

$$G_\lambda(x,y) \simeq d_*(x,y)\, N\left(x, \frac{1}{d_*(x,y)}\right) \quad \text{as } d_*(x,y) \to 0.$$

(2) *If X is non-compact then*

$$G_\lambda(x,y) \simeq \frac{1}{d_*(x,y)} N\left(x, \frac{1}{d_*(x,y)}\right) \quad \text{as } d_*(x,y) \to \infty\,.$$

The *potential operator* R associated with the semigroup $\{P^t\}_{t>0}$ is defined on the set of non-negative Borel measurable functions f as

$$Rf(x) = \lim_{\lambda \to 0} R_\lambda f(x).$$

Definition 5.3 (1) *The process $\{X_t\}_{t>0}$ and the semigroup $\{P^t\}_{t>0}$ are called* transient *if the potential Rf is a bounded function whenever f is bounded and has compact support.*
(2) *The* Green function *$G(x,y)$ is defined as* $G(x,y) = \lim_{\lambda \to 0} G_\lambda(x,y)$.

Using (5.1), we obtain

$$G(x,y) = \int_0^{1/d_*(x,y)} \frac{N(x,\tau)\,d\tau}{\tau^2} \le \infty. \tag{5.2}$$

Obviously, transience requires non-compactness of X, and when X is compact, $G(x,y) = \infty$ for all $x,y \in X$ (compare with Figure 1).

Theorem 5.4 *Suppose that (X,d) is non-compact. Then the following statements are equivalent.*

(1) *The semigroup $\{P^t\}_{t>0}$ is transient.*

(2) *$G(x,y) < \infty$ for some/all distinct $x,y \in X$.*

(3) *For some/all $x \in X$,*

$$\int_0^1 \frac{N(x,\tau)\, d\tau}{\tau^2} < \infty\,.$$

Proof. The equivalence of statements (2) and (3) follows from the very definition of the Green function. To prove the equivalence (1) $\Leftrightarrow$ (2), set $f = \mathbf{1}_B$, where $B = B_r^*(a)$ with $a \in X$ and $r \in \Lambda_*(a)$, as defined in (3.1).

Consider the potential $R(x,B) := Rf(x)$. By the Maximum Principle,

$$\sup_{x\in X} R(x,B) = \sup_{x\in B} R(x,B).$$

Let $B' = B_{r'}(a)$ be the predecessor ball of B in the sense of Definition 2.1. It exists by non-compactness of X. Write

$$\begin{aligned} R(x,B) &= \int_B G(x,y)\, d\mu(y) \\ &= \int_B d\mu(y) \int_0^{1/d_*(x,y)} \frac{N(x,\tau)\, d\tau}{\tau^2} = \int_{(0\,,\,r')} dV(x,s) \int_0^{1/s} \frac{N(x,\tau)\, d\tau}{\tau^2}, \end{aligned}$$

where $dV(x,s)$ refers to Lebesgue-Stieltjes integration with respect to the intrinsic volume function (1.7) at x. Changing the order of integration we obtain

$$\begin{aligned} R(x,B) &= \int_0^\infty \frac{N(x,\tau)}{\tau^2} \left(\int_{(0\,,\,r')} \mathbf{1}_{(0\,,\,1/s]}(\tau)\, dV(x,s) \right) d\tau \\ &= \int_0^\infty \frac{N(x,\tau)}{\tau^2} \left(\int_{(0\,,\,r')\cap(0\,,\,1/\tau]} dV(x,s) \right) d\tau. \end{aligned}$$

By Definition 3.2 , $N(x,\tau)V(x,\frac{1}{\tau}) = 1$ for all $\tau > 0$. Hence

$$\begin{aligned} R(x,B) &= \int_0^{1/r'} \frac{N(x,\tau)}{\tau^2} V(x,r)\, d\tau + \int_{1/r'}^\infty \frac{N(x,\tau)}{\tau^2} V(x,\frac{1}{\tau})\, d\tau \\ &= V(x,r) \int_0^{1/r'} \frac{N(x,\tau)}{\tau^2}\, d\tau + r'. \end{aligned}$$

Observe that $r' = d_*(x,z)$ for any $x \in B$ and any $z \in B' \setminus B$. Moreover $N(x,\tau) = N(y,\tau)$ for any $x,y \in B$ and $\tau \le 1/r'$. It follows that for any $x \in B$ and any $z \in B' \setminus B$,

$$R(x,B) = r' + \mu(B)G(a,z).$$

The last identity yields the equivalence of (1) and (2). ∎

When X is discrete and satisfies the conditions (*i*)–(*iii*) of the Introduction, i.e. the group of isometries acts transitively, then X can be seen as a locally finite group. If μ is the Haar measure, and we consider the discrete time processes, then the transience criterion (3) of the last theorem translates into the the general sufficient transience condition of [27]. Thus, the latter is also the necessary one.

Corollary 5.5 *Let (X,d) be non-discrete. Assume that the semigroup $\{P^t\}$ is transient and that $G(x,x) = \infty$ for some (equivalently, all) x. Then*

(1) *For any given $\lambda > 0$, as $d_*(x,y) \to 0$,*

$$G_\lambda(x,y) \sim G(x,y).$$

(2) *Assume in addition that the function* $\tau \mapsto N(x,\tau)$ *is doubling. Then, as* $d_*(x,y) \to 0$,

$$G(x,y) \simeq d_*(x,y)\, N\left(x, \frac{1}{d_*(x,y)}\right).$$

Consider the transient case. Fix $x \in X$, set $d_*(x,y) = r$ and $\varphi(\tau) = \tau N(x, 1/\tau)$, and write

$$G(x,y) = \int_r^\infty \varphi(\tau)\, \frac{d\tau}{\tau}.$$

A typical result in Karamata theory reads as follows.

$$\int_r^\infty \varphi(\tau)\, \frac{d\tau}{\tau} \simeq \varphi(r) \quad \text{as} \quad r \to \infty$$

if and only if the function F satisfies certain Tauberian conditions, see Bingham, Goldie and Teugels [9, Corollary 2.6.4]. In our setting, this means that

$$G(x,y) \simeq d_*(x,y)\, N\left(x, \frac{1}{d_*(x,y)}\right), \quad \text{as} \quad d_*(x,y) \to \infty$$

if and only if the function $\tau \mapsto \tau N(x, 1/\tau)$ satisfies the above mentioned Tauberian conditions. A simple sufficient condition for the last property to hold is given in the next theorem.

Theorem 5.6 *Assume that there exist constants* $0 < \varepsilon'' < \varepsilon' < \varepsilon < 1$ *such that*

$$\varepsilon'' \le \frac{N(x,\varepsilon\tau)}{N(x,\tau)} \le \varepsilon' \tag{5.3}$$

for some $x \in X$ *and all* $0 < \tau \le 1$. *Then the semigroup* $\{P^t\}_{t>0}$ *is transient and, for all* $y \in X$ *such that* $d_*(x,y) \ge 1$,

$$G(x,y) \simeq d_*(x,y) N\left(x, \frac{1}{d_*(x,y)}\right).$$

Proof. For any $0 < \eta \le 1$ we have

$$\int_0^\eta \frac{N(x,\tau)}{\tau^2}\, d\tau = \sum_{k=0}^\infty \int_{\varepsilon^{k+1}\eta}^{\varepsilon^k \eta} \frac{N(x,\tau)}{\tau^2}\, d\tau = \sum_{k=0}^\infty \int_{\varepsilon\eta}^{\eta} \frac{N(x,\varepsilon^k\tau)}{\tau^2} \varepsilon^{-k}\, d\tau.$$

Using the upper bound in (5.3), we obtain

$$\begin{aligned}\int_0^\eta \frac{N(x,\tau)}{\tau^2}\, d\tau \ &\le\ \sum_{k=0}^\infty \int_{\varepsilon\eta}^{\eta} \frac{N(x,\tau)}{\tau^2} (\varepsilon')^k \varepsilon^{-k}\, d\tau \le \frac{1-\varepsilon}{1-\varepsilon'/\varepsilon} \eta N(x,\eta) \frac{1}{(\varepsilon\eta)^2} \\ &=\ \frac{1-\varepsilon}{\varepsilon(\varepsilon-\varepsilon')} \frac{N(x,\eta)}{\eta} = c_2 \frac{N(x,\eta)}{\eta}.\end{aligned}$$

Analogously, using the lower bound in (5.3) we get

$$\begin{aligned}\int_0^\eta \frac{N(x,\tau)}{\tau^2}\, d\tau \ &\ge\ \int_{\varepsilon\eta}^\eta \frac{N(x,\tau)}{\tau^2}\, d\tau \ge \frac{1-\varepsilon}{\varepsilon} \frac{N(x,\varepsilon\eta)}{\eta} \\ &\ge\ \frac{\varepsilon''(1-\varepsilon)}{\varepsilon} \frac{N(x,\eta)}{\eta} = c_1 \frac{N(x,\eta)}{\eta}.\end{aligned}$$

Setting $\eta = 1/d_*(x,y)$ and using (5.2), we come to the desired conclusion. ■

Example 5.7 This is a continuation of Example 4.11: $X = \mathbb{Q}_p$ is the field of p-adic numbers with standard distance $d(x,y) = \|x-y\|_p$ and with modified distance $d_*(x,y) = p^{-\alpha}\|x-y\|_p^{\alpha}$, $\alpha > 0$. The spectral distribution function $N(x,\tau)$ does not depend on x, whence we denote it $N(\tau)$. It satisfies

$$p^{-1}\tau^{1/\alpha} \le N(\tau) \le \tau^{1/\alpha}.$$

By Theorem 5.4, the semigroup $\{P^t\}_{t>0}$ is transient if and only if $\alpha < 1$. Moreover, for all x, y,

$$G(x,y) \simeq \|x-y\|_p^{-1+\alpha}.$$

Note that in this example (and in many others) the condition (5.3) is in fact equivalent to transience. Indeed, for any fixed $0 < \varepsilon < 1$ and all $\tau > 0$,

$$\varepsilon^{1/\alpha} \le \frac{N(\varepsilon\tau)}{N(\tau)} \le p\,\varepsilon^{1/\alpha}.$$

When $\alpha < 1$, we can choose $0 < \varepsilon < p^{-\alpha/(1-\alpha)}$, $\varepsilon' = p\,\varepsilon^{1/\alpha}$ and $\varepsilon'' = \varepsilon^{1/\alpha}$ to show that the condition (5.3) holds. When $\alpha \ge 1$,

$$\varepsilon \le \varepsilon^{1/\alpha} \le \frac{N(\varepsilon\tau)}{N(\tau)},$$

whence the condition (5.3) does not hold.

6 Moments of the Markov process

Let $\{\mathcal{X}_t\}$ be the Markov process associated with the semigroup $\{P_t\}_{t>0}$. For any $\gamma > 0$, the moment of order γ of the process is defined as

$$M_\gamma(x,t) = \mathbb{E}_x\Big(d_*(x,\mathcal{X}_t)^\gamma\Big),$$

where $\mathbb{E}_x$ is expectation with respect to the probability measure on the trajectory space of $\{\mathcal{X}_t\}$ that governs the process starting at x. In terms of the transition function $p(t,x,y)$, that moment is given by

$$M_\gamma(x,t) = \int_X d_*(x,y)^\gamma p(t,x,y)\,d\mu(y). \tag{6.1}$$

The aim of this section is to estimate $M_\gamma(x,t)$ as a function of t and γ. For the main result, we shall need two lemmas. Recall the intrinsic volume function (1.7). We also need its average moment function of order γ at x,

$$R_\gamma(x,\tau) = \frac{1}{V(x,\tau)}\int_{(0,\tau]} r^\gamma\,dV(x,r).$$

Lemma 6.1 *For all $x \in X$, $t > 0$ and $\gamma > 0$,*

$$M_\gamma(x,t) = \int_0^\infty R_\gamma\left(x,\frac{1}{\tau}\right) t\,e^{-\tau t}\,d\tau = \int_0^\infty R_\gamma\left(x,\frac{t}{\tau}\right) e^{-\tau}\,d\tau.$$

Proof. Using the equations (6.1) and (3.3), as well as the Definition 3.2 of the spectral distribution function in terms of the volume function, we obtain

$$\begin{aligned}
M_\gamma(x,t) &= \int_X d_*(x,y)^\gamma\,p(t,x,y)\,d\mu(y)\\
&= \int_{(0\,,\,\infty)} r^\gamma\left(t\int_0^{1/r} N(x,\tau)\,e^{-\tau t}\,d\tau\right)dV(x,r)\\
&= \int_0^\infty\left(\int_{(0\,,\,1/\tau)}\frac{r^\gamma}{V(x,1/\tau)}\,dV(x,r)\right)t\,e^{-\tau t}\,d\tau = \int_0^\infty R_\gamma\left(x,\frac{1}{\tau}\right)t\,e^{-\tau t}\,d\tau.
\end{aligned}$$

In the 3rd identity, we have used Fubini's theorem. ■

The volume function $r \mapsto V(x,r)$ non-decreasing. In view of its relation with the spectral distribution function (see Definition 3.2) it is a step function whose shape can be understood from figures 1 – 3. The function varies from 0 to $\mu(X)$. In the compact case, $V(x,r) = \mu(X)$ for all $r \geq r^*_{\max} = r^*_{\max}(x)$, the largest value in $\Lambda_*(x)$; compare with (2.2) and (3.1). When x is isolated, $V(x,r) = \mu\{x\}$ for all $0 \leq r < r^*_0 = r^*_0(x)$, the smallest positive value in $\Lambda_*(x)$.

Lemma 6.2 *For any given $x \in X$ and $\gamma > 0$, the following properties hold.*

(a) *The function $\tau \mapsto R_\gamma(x,\tau)$ is non-decreasing.*
If X is compact $R_\gamma(x,\tau) = R_\gamma\big(x, r^*_{\max}(x)\big)$ for all $\tau \geq r^*_{\max}(x)$.
If X is discrete and infinite, $R_\gamma(x,\tau) = R_\gamma\big(x, r^*_0(x)\big)$ for all $0 < \tau \leq r^*_0(x)$.

(b) *For all $\tau > 0$, we have*

$$R_\gamma(x,\tau) \leq \tau^\gamma$$

and, if the volume function $r \mapsto V(x,r)$ satisfies the reverse doubling property, then there exists a constant $c > 0$, such that

$$R_\gamma(x,\tau) \geq c\,\tau^\gamma \tag{6.2}$$

for all $\tau > 0$. In the non-discrete compact case, if the volume function satisfies the reverse doubling property at zero, (6.2) holds for all $0 < \tau < r^*_{\max}(x)$. In the discrete infinite case, if the volume function satisfies the reverse doubling property at infinity, (6.2) holds for all $\tau > r^*_0(x)$.

Proof. For the first part of (a), we integrate by parts:

$$R_\gamma(x,\tau) = \frac{1}{V(x,\tau)}\left(\tau^\gamma\, V(x,\tau) - \int_{(0\,,\,\tau]} V(x,s)\, ds^\gamma\right) = \int_{(0\,,\,\tau]} \left(1 - \frac{V(x,s)}{V(x,\tau)}\right) ds^\gamma,$$

whence $\tau \mapsto \mathcal{R}_\gamma\,(x,\tau)$ is non-decreasing.

The second part (a) is straightforward.

Regarding (b), the general upper bound on $R_\gamma(x,\tau)$ is obvious. If the volume function satisfies the reverse doubling property, then in the respective range,

$$\begin{aligned} R_\gamma(x,\tau) \ &\geq\ \frac{1}{V(x,\tau)}(\delta\tau)^\gamma\Big(V(x,\tau) - V(x,\delta\tau)\Big) \\ &=\ (\delta\tau)^\gamma\left(1 - \frac{V(x,\delta\tau)}{V(x,\tau)}\right) \geq \delta^\gamma(1-\kappa)\tau^\gamma = c\,\tau^\gamma \end{aligned}$$

for suitable constants $0 < \kappa, c < 1$. ■

Now, in order to estimate the moment function $t \mapsto M_\gamma(x,t)$, we need to estimate a Laplace-type integral as given by the formula of Lemma 6.1. We will treat such estimates in the two technical Propositions 6.6 and 6.7 at the end of this section. Before that, in the next three theorems, we anticipate the statements of the results regarding the moment function.

Theorem 6.3 *Assume that (X,d) is non-compact and has no isolated points. Then the following properties hold.*

(1) *For all $x \in X$, $t > 0$ and $0 < \gamma < 1$,*

$$M_\gamma(x,t) \leq \frac{t^\gamma}{1-\gamma}.$$

(2) *If for some* $x \in X$*, the volume function satisfies the reverse doubling property, then for any* $0 < \gamma < 1$,

$$M_\gamma(x,t) \geq \frac{c}{1-\gamma} t^\gamma,$$

for all $x, t > 0$ *and some* $c > 0$*. Moreover,*

$$M_\gamma(z,t) = \infty,$$

for all z, $t > 0$ *and* $\gamma \geq 1$.

Theorem 6.4 *Assume that* (X,d) *is discrete and infinite. Then the following properties hold.*

(a) *For all* x, $t > 0$ *and* $0 < \gamma < 1$,

$$M_\gamma(x,t) \leq \frac{C}{1-\gamma} \min\{t, t^\gamma\}$$

for some $C > 0$.

(b) *If for some (equivalently, all)* $x \in X$ *the volume function satisfies the reverse doubling property at infinity, then for any* $0 < \gamma < 1$,

$$M_\gamma(z,t) \geq \frac{c}{1-\gamma} \min\{t, t^\gamma\}$$

for all z , $t > 0$ *and for some* $c > 0$*. Moreover,*

$$M_\gamma(z,t) = \infty$$

for all z, $t > 0$ *and all* $\gamma \geq 1$.

Assume now that (X,d) is compact and let D be its d_*-diameter. By Lemmas 6.1 and 6.2, for all $x \in X$, $\gamma > 0$ and $t > 0$,

$$M_\gamma(x,t) \leq R_\gamma(x,D) \leq D^\gamma,$$

whence we study the behavior of the moment function $t \mapsto M_\gamma(x,t)$ at zero.

Theorem 6.5 *Assume that* (X,d) *is non-discrete and compact. Then the following properties hold.*

(1) *There exists a constant* $C > 0$ *such that*

$$M_\gamma(x,t) \leq \begin{cases} C\,t & \text{if} \quad \gamma > 1, \\ C\,t\,\left(\log\frac{1}{t} + 1\right) & \text{if} \quad \gamma = 1, \\ C\,t^\gamma & \text{if} \quad \gamma < 1, \end{cases}$$

holds for all x *and all* $0 < t \leq 1$.

(2) *If for some* $x \in X$ *the volume function satisfies the reverse doubling property at zero, then there exists a constant* $c > 0$ *such that*

$$M_\gamma(z,t) \geq \begin{cases} c\,t & \text{if} \quad \gamma > 1, \\ c\,t\,\left(\log\frac{1}{t} + 1\right) & \text{if} \quad \gamma = 1, \\ c\,t^\gamma & \text{if} \quad \gamma < 1 \end{cases}$$

holds for all z *and all* $0 < t \leq 1$.

We now provide the technical details regarding the Laplace-type estimates that imply Theorems 6.3, 6.4 and 6.5. In the following two propositions, M and R will always be two non-negative, non-decreasing functions related by the Laplace-type integral

$$M(t) = \int_0^\infty R\left(\frac{t}{\tau}\right) e^{-\tau}\, d\tau\,.$$

Proposition 6.6 *Let $\gamma > 0$ be given.*

(1) *Assume that*

$$A s^\gamma \geq R(s)\,, \quad \text{or that respectively} \quad R(s) \geq B\, s^\gamma \tag{6.3}$$

for some $A > 0$ (resp. $B > 0$) and all $s > 0$. Then the inequality

$$\frac{A\, t^\gamma}{1-\gamma} \geq M(t)\,, \quad \text{respectively} \quad M(t) \geq \frac{B\, t^\gamma}{(1-\gamma)\, e}$$

holds for all $0 < \gamma < 1$ and all $t > 0$.

(2) *Assume that there is $t_0 > 0$ such that $R(s) = 0$ for all $0 < s < t_0$. Assume also that one of the respective inequalities of (6.3) holds for all $s > t_0$. Then*

$$M(t) \leq \frac{c}{1-\gamma} \min\left\{\frac{t}{t_0}, \left(\frac{t}{t_0}\right)^\gamma\right\}, \quad \text{respectively} \quad M(t) \geq \frac{c'}{1-\gamma} \min\left\{\frac{t}{t_0}, \left(\frac{t}{t_0}\right)^\gamma\right\},$$

for all $0 < \gamma < 1$, all $t > 0$ and some constants $c, c' > 0$.

(3) *The assumption $\gamma \geq 1$ and the lower bound $R(s) \geq B\, s^\gamma$ imply that $M(t) = \infty$ for all $t > 0$.*

Proof. It is known that for $0 < \gamma < 1$ the Gamma-function satisfies

$$\frac{1}{(1-\gamma)\, e} < \Gamma(1-\gamma) < \frac{1}{1-\gamma},$$

whence by monotonicity of the Laplace-type integral the first claim follows.

To prove the second statement, we write

$$M(t) = \int_{\{t/s \geq t_0\}} R\left(\frac{t}{s}\right) e^{-s}\, ds.$$

First assume that $R(\tau) \leq A\, s^\gamma$ for all $0 < s < \infty$. Then we obtain

$$\begin{aligned} M(t) &\leq A \int_{\{t/s \geq t_0\}} \left(\frac{t}{s}\right)^\gamma e^{-s}\, ds = A\, t^\gamma \int_{\{s \leq t/t_0\}} s^{-\gamma} e^{-s}\, ds \\ &\leq A\, t^\gamma \int_0^{t/t_0} s^{-\gamma}\, ds = \left(\frac{t}{t_0}\right) \frac{A t_0^{-\gamma}}{1-\gamma}, \quad \text{and} \end{aligned}$$

$$M(t) \leq A\, t^\gamma \int_0^\infty s^{-\gamma} e^{-s}\, ds \leq \frac{A\, t^\gamma}{1-\gamma} = \left(\frac{t}{t_0}\right)^\gamma \frac{A\, t_0^\gamma}{1-\gamma}.$$

It follows that

$$M(t) \leq \frac{A \max\left\{t_0, t_0^{-1}\right\}}{1-\gamma} \min\left\{\frac{t}{t_0}, \left(\frac{t}{t_0}\right)^\gamma\right\}.$$

Second, assume that $R(s) \geq B\,s^\gamma$, for all $s \geq t_0$. Then for $t/t_0 \geq 1$

$$M(t) \geq B\,t^\gamma \int_0^{t/t_0} s^{-\gamma} e^{-s}\,ds \geq \frac{B\,t^\gamma}{e} \int_0^1 s^{-\gamma}\,ds = \frac{B\,t^\gamma}{(1-\gamma)e} = \frac{B\,t_0^\gamma}{(1-\gamma)\,e} \left(\frac{t}{t_0}\right)^\gamma .$$

When $t/t_0 \leq 1$ we get

$$M(t) \geq B\,t^\gamma \int_0^{t/t_0} s^{-\gamma} e^{-s}\,ds \geq \frac{B\,t^\gamma}{e} \int_0^{t/t_0} s^{-\gamma}\,ds = \frac{B\,t^\gamma}{(1-\gamma)\,e} \left(\frac{t}{t_0}\right)^{1-\gamma} = \frac{B\,t_0^\gamma}{(1-\gamma)\,e} \left(\frac{t}{t_0}\right) .$$

It follows that

$$M(t) \geq \frac{B\,t_0^\gamma}{(1-\gamma)\,e} \min\left\{\frac{t}{t_0}, \left(\frac{t}{t_0}\right)^\gamma\right\} \geq \frac{B \min\{t_0, 1\}}{(1-\gamma)\,e} \min\left\{\frac{t}{t_0}, \left(\frac{t}{t_0}\right)^\gamma\right\} .$$

This proves the second claim. For the third claim observe that that if $R(s) \geq B\,s^\gamma$ for all $s \geq t_0$ and $\gamma \geq 1$,

$$M(t) \geq B\,t^\gamma \int_0^{t/t_0} s^{-\gamma} e^{-s}\,ds = \infty$$

for all $t > 0$. ∎

Proposition 6.7 *Assume that there is* $t_0 > 0$ *such that* $R(s) = R(t_0)$ *for all* $s \geq t_0$. *Assume also that one of the respective inequalities in (6.3) holds for all* $0 < s \leq t_0$. *Then*

$$M(t) \quad \leq \quad \begin{cases} c_1 \frac{t}{t_0} & \textit{if} \quad \gamma > 1, \\ c_2\,t\left(\log \frac{t_0}{t} + 1\right) & \textit{if} \quad \gamma = 1, \\ c_3 \left(\frac{t}{t_0}\right)^\gamma & \textit{if} \quad \gamma < 1. \end{cases}$$

$$\textit{respectively,} \quad M(t) \quad \geq \quad \begin{cases} c_1' \frac{t}{t_0} & \textit{if} \quad \gamma > 1, \\ c_2'\,t\left(\log \frac{t_0}{t} + 1\right) & \textit{if} \quad \gamma = 1, \\ c_3' \left(\frac{t}{t_0}\right)^\gamma & \textit{if} \quad \gamma < 1. \end{cases}$$

for all $0 < t \leq t_0$ *and some positive constants* $c_1\,, c_1'\,, c_2\,, c_2'\,, c_3\,, c_3'$.

Proof. Let $\gamma > 1$ and $0 < t < t_0$. According to our assumption

$$M(t) = \int_{\{t/s \leq t_0\}} R\left(\frac{t}{s}\right) e^{-s}\,ds + R(t_0)\left(1 - e^{-t/t_0}\right) .$$

Observe that for $0 < t < t_0$,

$$\frac{t}{2t_0} \leq \left(1 - e^{-t/t_0}\right) \leq \frac{t}{t_0} .$$

First, if $R(s) \leq A\,s^\gamma$ for all $0 < s < t_0$, then

$$\begin{aligned} M(t) &\leq A\,t^\gamma \int_{t/t_0}^\infty s^{-\gamma} e^{-s}\,ds + \frac{R(t_0)t}{t_0} \leq A\,s^\gamma \int_{t/t_0}^\infty s^{-\gamma}\,ds + \frac{R(t_0)t}{t_0} \\ &\leq \frac{A\,t^\gamma}{\gamma - 1} \left(\frac{t}{t_0}\right)^{1-\gamma} + \frac{R(t_0)t}{t_0} = \frac{t}{t_0}\left(R(t_0) + \frac{A\,t_0^\gamma}{\gamma - 1}\right) . \end{aligned}$$

Second, if $R(s) \geq B\,s^\gamma$, for all $0 < s < t_0$, then

$$M(t) \geq \frac{R(t_0)}{2} \frac{t}{t_0} .$$

Assume that $0<\gamma<1$ and $0<t<t_0$. Again first, if $R(s)\le A\,s^\gamma$ for all $0<\tau<t_0$, then

$$\begin{aligned}
M(t) &\le At^\gamma\int_{t/t_0}^\infty s^{-\gamma}e^{-s}\,ds+\frac{R(t_0)t}{t_0}\le A\,t^\gamma\,\Gamma(1-\gamma)+R(t_0)\frac{t}{t_0}\\
&\le \frac{A\,t^\gamma}{1-\gamma}+R(t_0)\frac{t}{t_0}=\frac{A\,t_0^\gamma}{1-\gamma}\left(\frac{t}{t_0}\right)^\gamma+R(t_0)\frac{t}{t_0}\\
&\le \left(\frac{t}{t_0}\right)^\gamma\left(\frac{A\,T^\gamma}{1-\gamma}+R(t_0)\right).
\end{aligned}$$

Second, once more, when $R(s)\ge B\,s^\gamma$, for all $0<s<T$, then

$$M(t)\ge B\,t^\gamma\int_{t/t_0}^\infty s^{-\gamma}e^{-s}\,ds\ge B\,t^\gamma\int_1^\infty s^{-\gamma}e^{-s}\,ds\ge\left(\frac{t}{t_0}\right)^\gamma\left(\frac{B\,\min\{t_0,1\}}{e^2}\right).$$

Finally, assume that $\gamma=1$ and $0<t<t_0$. First, if $R(s)\le A\,s^\gamma$ for all $0<\tau<t_0$, then

$$\begin{aligned}
M(t) &\le At\int_{t/T}^\infty s^{-1}e^{-s}ds+\frac{R(T)t}{T}\\
&= At\left(\int_1^\infty s^{-1}e^{-s}\,ds+\int_{t/t_0}^1 s^{-1}e^{-s}\,ds\right)+\frac{R(t_0)t}{t_0}\\
&\le A\,t\left(\int_1^\infty\frac{ds}{s^2}+\int_{t/t_0}^1\frac{ds}{s}\right)+\frac{R(t_0)t}{t_0}=\left(A+\frac{R(t_0)}{t_0}\right)t\left(\log\frac{t_0}{t}+1\right).
\end{aligned}$$

And at last, if $R(s)\ge B\,s^\gamma$ for all $0<\tau<t_0$, then

$$\begin{aligned}
M(t) &\ge B\,t\int_{t/t_0}^\infty s^{-1}e^{-s}\,ds+\frac{R(t_0)t}{2t_0}\\
&\ge \frac{B\,t}{e}\int_{t/t_0}^1\frac{ds}{s}+\frac{R(t_0)t}{2t_0}=\frac{B\,t}{e}\log\frac{t_0}{t}+\frac{R(t_0)t}{2t_0}\\
&= \frac{B\,t}{e}\left(\log\frac{t_0}{t}+\frac{R(t_o)e}{2B\,t_0}\right)\ge\min\left\{\frac{R(t_0)}{2t_0},\frac{B}{e}\right\}t\left(\log\frac{t_0}{t}+1\right).
\end{aligned}$$

The proof is finished. ∎

Theorems 6.3, 6.4 and 6.5 follow.

7 The Markov generator and its spectrum

It is known that any strongly continuous semigroup $\{P_t\}_{t>0}$ in a Banach space B has the infinitesimal generator $(-\mathcal{L},\mathsf{Dom}_\mathcal{L})$, whose domain is

$$\mathsf{Dom}_\mathcal{L}=\left\{f\in B:\lim_{t\to0}\frac{f-P_tf}{t}\text{ exists strongly}\right\},$$

and, for $f\in\mathsf{Dom}_\mathcal{L}$,

$$\mathcal{L}f=\lim_{t\to0}\frac{f-P_tf}{t}.$$

In general, $(\mathcal{L},\mathsf{Dom}_\mathcal{L})$ is an unbounded densely defined closed operator in B. Moreover, if B is a Hilbert space and the operators P_t are symmetric then $(\mathcal{L},\mathsf{Dom}_\mathcal{L})$ is a self-adjoint operator and $P_t=\exp(-t\mathcal{L})$. The purpose of this section is to study the infinitesimal generator $(-\mathcal{L},\mathsf{Dom}_\mathcal{L})$

of our symmetric Markov semigroup $\{P^t\}_{t>0}$ given by (1.4). We will refer to $(\mathcal{L}, \mathsf{Dom}_{\mathcal{L}})$ as the *Laplace operator* (shortly, *Laplacian*) associated to the symmetric Markov semigroup $\{P^t\}_{t>0}$. This is a non-negative definite self-adjoint operator.

Let us introduce the function

$$J_t(x,y) := \int_0^{1/d_*(x,y)} N(x,\tau)\exp(-\tau t)\,d\tau,$$

defined for all $x, y \in X$ and all $t \geq 0$. In particular, for $t = 0$ we set

$$J(x,y) := J_0(x,y) = \int_0^{1/d_*(x,y)} N(x,\tau)\,d\tau. \tag{7.1}$$

By construction, and in particular by (2.4), the functions $J_t(x,y)$ and $J(x,y)$ are symmetric and finite on the set $\{(x,y) \in X^2 : x \neq y\}$, and $J(x,x) = \infty$ for all x, whereas $J_t(x,x)$ may be finite for some $t > 0$ and $x \in X$. We shall see that $J(x,y)$ plays the role of the jump-kernel of the Dirichlet form associated with our Laplacian.

Yet one more observation is in order: for all x, y, $\lambda > 0$,

$$J(x,y) > \lambda^2 G_\lambda(x,y),$$

and in the non-compact case, for any fixed $\lambda > 0$,

$$J(x,y) \sim \lambda^2 G_\lambda(x,y) \quad \text{as} \;\; d_*(x,y) \to \infty$$

by Theorem 5.1.

Lemma 7.1 *The following properties hold.*

(a) *For all $x, y \in X$,*

$$\frac{1}{2d_*(x,y)} N\left(y, \frac{1}{2d_*(x,y)}\right) \leq J(x,y) \leq \frac{1}{d_*(x,y)} N\left(y, \frac{1}{d_*(x,y)}\right).$$

(b) *Let $B = B_r^*(a)$ be a fixed ball of radius $r > 0$. Then*

$$\int_{X\setminus B} J(x,y)\,d\mu(y) \leq \frac{1}{r} \quad \text{for any } \; x \in B, \text{ and}$$

$$\int_{X\setminus B} \left(\int_B J(x,y)\,d\mu(y)\right)^2 d\mu(x) \leq \frac{V(a,r)}{r^2}.$$

Proof. Statement (a) is immediate from the definition of the function $J(x,y)$ and the fact that the function $\tau \to N(x,\tau)$ is non-decreasing.

For (b), we may of course assume that $B \subsetneq X$. Recall the structure of the sets $\Lambda(a)$ and $\Lambda_*(a)$ from (2.2) and (3.1). Let once more B' be the predecessor ball of B and r' its radius (see Definition 2.1). It exists because $B \subsetneq X$. All $x \in B$ and $y \in X \setminus B$ satisfy $d_*(x,y) = d_*(a,y) \geq r' > r$, whence $J(x,y) = J(a,y)$. With this observation in mind we compute for any $z \in B' \setminus B$

$$\begin{aligned}
\int_{X\setminus B} J(x,y)\,d\mu(y) &= \int_{(r',\infty)} dV(a,s) \int_0^{1/s} \frac{d\tau}{V(a,1/\tau)} \\
&= \int_0^{1/r'} d\tau \int_{(r',1/\tau]} \frac{dV(a,s)}{V(a,1/\tau)} = \int_0^{1/r'} \frac{V(a,1/\tau) - V(a,r')}{V(a,1/\tau)}\,d\tau \\
&= \frac{1}{r'} - V(a,r') \int_0^{1/r'} \frac{d\tau}{V(a,1/\tau)} = \frac{1}{r'} - V(a,r')J(a,z),
\end{aligned} \tag{7.2}$$

Since $r' > r$, this proves the first identity of (b).

To prove the second identity of (b), we observe that when $y \in B$ and $x \in X \setminus B$,

$$J(x,y) = J(x,a) \leq \frac{1}{d_*(x,a)} N\left(a, \frac{1}{d_*(x,a)}\right). \tag{7.3}$$

Using the inequality 7.3 we write

$$\begin{aligned}
\int_{X\setminus B}\left(\int_B J(x,y)d\mu(y)\right)^2 d\mu(x) &\leq V(a,r)^2 \int_{X\setminus B}\left(\frac{1}{d_*(x,a)}N\Big(a, \frac{1}{d_*(x,a)}\Big)\right)^2 d\mu(x) \\
&\leq \left(\frac{V(a,r)}{r}\right)^2 \int_{X\setminus B}\frac{d\mu(x)}{V\big(a, d_*(x,a)\big)^2}.
\end{aligned}$$

Write $\Lambda_*(a) = \{0\} \cup \{r_k\}$, where k varies as in (2.2), but refers to the intrinsic metric d_*. Then

$$\begin{aligned}
\int_{X\setminus B}\frac{d\mu(x)}{V\big(a, d_*(x,a)\big)^2} &= \sum_{r_k \geq r'} \frac{V(a,r_k) - V(a,r_{k-1})}{\big(V(a,r_k)\big)^2} \\
&< \sum_{r_k > r} \frac{V(a,r_k) - V(a,r_{k-1})}{V(a,r_k)V(a,r_{k-1})} = \frac{1}{V(a,r)},
\end{aligned}$$

which concludes the proof. ■

As a set of test functions, we introduce the linear space $\mathcal{D}$ of all real valued, locally constant and compactly supported functions on X. Evidently $\mathcal{D}$ is dense in all Banach spaces $L^p = L^p(X,\mu)$, $1 \leq p < \infty$, as well as in $C_0(X)$, the space of all continuous functions vanishing at infinity.

The following theorem shows that the Laplacian arises from a difference operator, and it leads to an explicit representation of the associated Dirichlet form.

Theorem 7.2 *Let $(\mathcal{L}, \mathsf{Dom}_{\mathcal{L}})$ be the Laplacian. Then $\mathcal{D} \subset \mathsf{Dom}_{\mathcal{L}}$ and for any $f \in \mathcal{D}$,*

$$\begin{aligned}
\mathcal{L}f(x) &= \int_X \big(f(x) - f(y)\big) J(x,y)\, d\mu(y), \quad \textit{and} \\
(\mathcal{L}f, f) &= \frac{1}{2}\iint_{X\times X}\big(f(x) - f(y)\big)^2 J(x,y)\, d\mu(x)\, d\mu(y).
\end{aligned}$$

Proof. For any continuous compactly supported function f we write

$$\frac{1}{t}\big(f(x) - P^t f(x)\big) = \int_X \big(f(x) - f(y)\big)\frac{p(t,x,y)}{t}\, d\mu(y).$$

Using the equation (3.3) we obtain

$$\begin{aligned}
\frac{1}{t}\big(f(x) - P^t f(x)\big) &= \int_X \big(f(x) - f(y)\big)\int_0^{1/d_*(x,y)} N(x,\tau)\exp(-\tau t)\, d\lambda\, d\mu(y) \\
&= \int_X \big(f(x) - f(y)\big) J_t(x,y)\, d\mu(y).
\end{aligned}$$

Now assume that f is a test function. Then f can be written as a finite linear combination of indicators $\mathbf{1}_{B_k}$ of non-intersecting balls B_k. Hence we can restrict our considerations to the case when $f = \mathbf{1}_B$, where $B = B_r^*(a)$ is a ball of radius $r > 0$. Under this assumption the function

$$U_t(x) = \int_X \big(f(x) - f(y)\big) J_t(x,y)\, d\mu(y)$$

has the form

$$U_t(x) = \begin{cases} \int_{X\setminus B} J_t(x,y)\,d\mu(y) & \text{if} \quad x \in B \\ -\int_B J_t(x,y)\,d\mu(y) & \text{if} \quad x \in X \setminus B. \end{cases}$$

As $t \to 0$ the function U_t converges pointwise to the similar function U,

$$U(x) = \begin{cases} \int_{X\setminus B} J(x,y)\,d\mu(y) & \text{if} \quad x \in B \\ -\int_B J(x,y)\,d\mu(y) & \text{if} \quad x \in X \setminus B. \end{cases} \tag{7.4}$$

By construction, $U_t \in L^2$ for all $t > 0$, and $U \in L^2$ by Lemma 7.1. Our above computations show that

$$\begin{aligned} \|U - U_t\|^2_{L^2(X,\mu)} &= \int_B \Big(\int_{X\setminus B} \big(J(x,y) - J_t(x,y)\big)\,d\mu(y) \Big)^2 d\mu(x) \\ &\quad + \int_{X\setminus B} \Big(\int_B \big(J(x,y) - J_t(x,y)\big)\,d\mu(y) \Big)^2 d\mu(x). \end{aligned}$$

Since $J(x,y) > J_t(x,y)$ and $J_t(x,y) \to J(x,y)$ monotonically as $t \to 0$, we can apply the Monotone Convergence Theorem to conclude that $U_t \to U$ in L^2. Thus

$$\mathcal{L}\mathbf{1}_B(x) = U(x) \tag{7.5}$$

and we see that indeed $\mathcal{D} \subset \mathsf{Dom}_{\mathcal{L}}$ and that the first formula holds for every test function.

We now prove the second formula. It is a general fact concerning any symmetric Markov operator which admits a transition density with respect to a reference measure that, in our specific setting, the following identity holds for any test function f:

$$\begin{aligned} \big(f - P^t f, f\big) &= \frac{1}{2} \iint_{X\times X} \big(f(x) - f(y)\big)^2 p(t,x,y)\,d\mu(x)\,d\mu(y)\,, \quad \text{whence} \\ \frac{1}{t}\big(f - P^t f, f\big) &= \frac{1}{2} \iint_{X\times X} \big(f(x) - f(y)\big)^2 J_t(x,y)\,d\mu(x)\,d\mu(y). \end{aligned}$$

We apply once more the Monotone Convergence Theorem and the fact that $\frac{1}{t}(f - P^t f) \to \mathcal{L}f$ to get the desired result. ■

Let $\mathcal{B}$ be the family of all closed balls in (X,d), or equivalently, in (X,d_*). For any $B \in \mathcal{B}$ we let $r^*(B)$ denote the d_*-diameter of B, or equivalently, its minimal radius, a number that belongs to the set Λ_* of (3.1). For any $B \in \mathcal{B}$, with $B \subsetneq X$ in the compact case, we define the function

$$f_B = \frac{1}{\mu(B)} \mathbf{1}_B - \frac{1}{\mu(B')} \mathbf{1}_{B'}, \tag{7.6}$$

where the ball B' is the the predecessor of B. By abuse of notation, we shall write $X' = X$. When X is compact, it also belongs to $\mathcal{B}$, and we set

$$f_X = \frac{1}{\mu(X)} \mathbf{1}_X\,.$$

Observe that $f_B \in \mathcal{D}$, its support is B' and, unless $B = X$,

$$\|f_B\|^2_{L^2} = \frac{1}{\mu(B)} - \frac{1}{\mu(B')},$$

while of course $\|f_X\|^2_{L^2} = 1/\mu(X)$ and $\mathcal{L}f_X = 0$ in the compact case. For $B, C \in \mathcal{B}$, we have $f_B \perp f_C$ if and only if $B' \neq C'$.

Theorem 7.3 (1) *The family of functions* $\{f_B : B \in \mathcal{B}\}$ *is a complete system in* L^2 *(i.e., its linear span is dense).*

(2) *Any function* f_B, $B \neq X$, *satisfies the equation*

$$\mathcal{L} f_B = \frac{1}{r^*(B')} f_B.$$

In particular, the symmetric operator $(\mathcal{L}, \mathcal{D})$ *has a complete system of eigenfunctions.*

Proof. (1) Assume that some function $f \in L^2$ is orthogonal to all functions f_B. Then for any $B \in \mathcal{B}$, $B \neq X$, the averages

$$\frac{1}{\mu(B)} \int_B f \, d\mu \quad \text{and} \quad \frac{1}{\mu(B')} \int_{B'} f \, d\mu$$

coincide. Since any two balls are contained in a common bigger ball, by induction the averages coincide over any two balls in $\mathcal{B}$. It follows that f must be a constant function. X is non-compact, then $f = 0$ because it belongs to $L^2(X, \mu)$ and $\mu(X) = \infty$ by assumption (2.1).

If X is compact, then $f = 0$ because it is orthogonal to the function $f_X = 1$. This proves (1).

(2) For a given ball $B \neq X$,

$$\mathcal{L} f_B(x) = \frac{1}{\mu(B)} \mathcal{L}\mathbf{1}_B(x) - \frac{1}{\mu(B')} \mathcal{L}\mathbf{1}_{B'}(x).$$

We distinguish the following three cases regarding the location of x.

Case1. $x \in X \setminus B'$. Then $f_B(x) = 0$.

By (7.4) and (7.5),

$$\mathcal{L} f_B(x) = -\frac{1}{\mu(B)} \int_B J(x,y) \, d\mu(y) + \frac{1}{\mu(B')} \int_{B'} J(x,y) \, d\mu(y).$$

Since $J(x,y) = J(x,a)$, for any $y, a \in B$,

$$\mathcal{L} f_B(x) = -\frac{\mu(B)}{\mu(B)} J(x,a) + \frac{\mu(B')}{\mu(B')} J(x,a) = 0.$$

Case2. $x \in B$. Then $f_B(x) = \frac{1}{\mu(B)} - \frac{1}{\mu(B')}$.

Again by (7.4) and (7.5),

$$\begin{aligned}
\mathcal{L} f_B(x) &= \frac{1}{\mu(B)} \int_{X \setminus B} J(x,y) \, d\mu(y) - \frac{1}{\mu(B')} \int_{X \setminus B'} J(x,y) \, d\mu(y) \\
&= \left(\frac{1}{\mu(B)} - \frac{1}{\mu(B')} \right) \int_{X \setminus B'} J(x,y) \, d\mu(y) + \frac{1}{\mu(B)} \int_{B' \setminus B} J(x,y) \, d\mu(y).
\end{aligned}$$

When $x \in B$ and $y \in X \setminus B'$, we have $d_*(x,y) = d_*(a,y)$ for any $a \in B$. Thus, by (7.2) , for any $z \in B' \setminus B$,

$$\begin{aligned}
\mathcal{L} f_B(x) &= \left(\frac{1}{\mu(B)} - \frac{1}{\mu(B')} \right) \left(\frac{1}{r^*(B')} - \mu(B') J(a,z) \right) + \frac{1}{\mu(B)} \left(\mu(B') - \mu(B) \right) J(a,z) \\
&= \frac{1}{r^*(B')} \left(\frac{1}{\mu(B)} - \frac{1}{\mu(B')} \right) = \frac{1}{r^*(B')} f_B(x).
\end{aligned}$$

Case 3. $x \in B' \setminus B$. Then $f_B(x) = -\frac{1}{\mu(B')}$. Once more by (7.4) and (7.5),

$$\mathcal{L} f_B(x) = -\frac{1}{\mu(B)} \int_B J(x,y)\, d\mu(y) - \frac{1}{\mu(B')} \int_{X \setminus B'} J(x,y)\, d\mu(y).$$

Our x belongs to a ball $C \neq B$ with $C' = B'$ (there is a finite number of such balls). We apply (7.2) and choosing an arbitrary $z \in B' \setminus C$, we obtain

$$\begin{aligned} \mathcal{L} f_B(x) &= -J(x,z) - \frac{1}{\mu(B')} \left(\frac{1}{r^*(B')} - \mu(B') J(x,z) \right) \\ &= \frac{1}{r^*(B')} \left(-\frac{1}{\mu(B')} \right) = \frac{1}{r^*(B')} f_B(x). \end{aligned}$$

This completes the proof of statement (2). ■

For $B \in \mathcal{B}$, we define the space

$$\mathcal{H}_B = \mathrm{Span}\{ f_C : C \in \mathcal{B},\ C' = B \}. \tag{7.7}$$

An easy computation shows that

$$\sum_{C \in \mathcal{B}\,:\, C' = B} \mu(C) f_C = 0. \tag{7.8}$$

On the other hand, if we select one $C_B \in \mathcal{B}$ with $C'_B = B$, then by another straightforward computation the functions $\{ f_C : C' = B\,,\ C \neq C_B \}$ are linearly independent. Combining these observations with Theorem 7.3, we now see that the spectrum of our Laplacian is pure point.

Corollary 7.4 (a) *The symmetric operator* $(\mathcal{L}, \mathcal{D})$ *is essentially self-adjoint.*

(b) *The spectrum of the self-adjoint operator* $(\mathcal{L}, \mathsf{Dom}_{\mathcal{L}})$ *coincides with the closure of the set*

$$\left\{ \frac{1}{r^*} : r^* \in \Lambda_*\,,\ r^* \neq 0 \right\} \cup \{0\} \qquad \text{with } \Lambda_* \text{ as in (3.1).}$$

(c) *The space* $L^2(X,\mu)$ *decomposes as a direct sum of finite-dimensional eigenspaces which are spanned by compactly supported functions:*

$$L^2(X,\mu) = \bigoplus_{B \in \mathcal{B}} \mathcal{H}_B.$$

It is now easy to prove our general subordination result. Recall the inverse exponential distribution σ_* of (1.5). Also recall from the introduction that our (d,μ,σ)-process and the associated Laplacian coincides with the (d_*,μ,σ_*)-process, which is the standard (d_*,μ)-process in our terminology.

Theorem 7.5 *Let* $\psi : [0\,,\,\infty) \to [0\,,\,\infty)$ *be an increasing bijection, and let* $(\mathcal{L}, \mathsf{Dom}_{\mathcal{L}})$ *be the Laplacian associated with the* (d,μ,σ)*-process.*

Then the self-adjoint operator $\big(\psi(\mathcal{L}), \mathsf{Dom}_{\psi(\mathcal{L})}\big)$ *is the Laplacian associated with the standard* (d_ψ,μ)*-process, where the new ultra-metric* d_ψ *on* X *is given by*

$$\frac{1}{d_\psi(x,y)} = \psi\left(\frac{1}{d_*(x,y)} \right).$$

(It is the intrinsic metric of the new process.)

Proof. We apply the L^2-decomposition of Corollary 7.4(c) and the fact that each space $\mathcal{H}_B \subset L^2$ is an eigenspace of the operator $(\mathcal{L}, \mathcal{D})$ corresponding to the eigenvalue $1/r^*(B')$. Let $\Pi_B : L^2 \to \mathcal{H}_B$ be the corresponding orthoprojector. Then, for any function $f \in \mathsf{Dom}_{\mathcal{L}}$,

$$f = \sum_{B \in \mathcal{B}} \Pi_{B'} f \quad \text{and} \quad \mathcal{L} f = \sum_{B \in \mathcal{B}} \frac{1}{r^*(B')} \Pi_B f. \tag{7.9}$$

(The sums are at most countable by the general theory.) Let $r_\psi(B')$ be the diameter of the ball B' with respect to the new ultra-metric d_ψ. By Spectral Theory, for any $f \in \mathsf{Dom}_{\psi(\mathcal{L})}$,

$$\psi(\mathcal{L}) f = \sum_{B \in \mathcal{B}} \psi\left(\frac{1}{r^*(B')}\right) \Pi_B f = \sum_{B \in \mathcal{B}} \frac{1}{d_\psi(B')} \Pi_B f. \tag{7.10}$$

Comparing the equations (7.9) and (7.10), we now conclude that the self-adjoint operator $\big(\psi(\mathcal{L}), \mathsf{Dom}_{\psi(\mathcal{L})}\big)$ coincides with the Laplacian of the standard (d_ψ, μ)-process on X. ■

Remark 7.6 Recall that a non-negative definite, self-adjoint operator $\mathcal{L}$ is a Laplacian if its semigroup $(e^{-t\mathcal{L}})_{t>0}$ is Markovian. In general, by Bochner's theorem, for any Laplace operator $\mathcal{L}$ the operator $\psi(\mathcal{L})$ is again a Laplace operator, provided that ψ is a Bernstein function. See for example SCHILLING, SONG AND VONDRAČEK [38]. Now, $\psi(\lambda) = \lambda^\alpha$ is a Bernstein function if and only if $0 < \alpha \leq 1$. Thus, for a general Laplacian, it is not true that $\mathcal{L}^\alpha$ with $\alpha > 1$ is a Laplace operator.

However, by Theorem 7.5, in our specific case of the isotropic semigroup $\{P^t\}_{t>0}$ plus associated Laplacian on an ultrametric space, as given by (1.4), the operator $\mathcal{L}^\alpha$ *is* a Laplace operator for any $\alpha > 0$.

Proposition 7.7 *Let (X, d) be a non-compact, proper ultra-metric space. Let $M \subset [0, \infty)$ be any closed set (unbounded, if X contains at least one non-isolated point) that accumulates at* 0. *Then the following holds.*

(a) *There exists a proper ultra-metric d' on X which generates the same topology as d and a Laplacian $(\mathcal{L}, Dom_{\mathcal{L}})$ on (X, d') such that*

$$\mathrm{Spec}(\mathcal{L}, \mathsf{Dom}_{\mathcal{L}}) = M.$$

(b) *Suppose in addition that there exists a partition of X into d-balls that consists of infinitely many non-singletons. Then the ultra-metric d' of* (a) *can be chosen such that the collections of d-balls and d'-balls coincide.*

Proof. The set

$$D = \{x \in (0, \infty) : x^{-1} \in M\} \cup \{0\}$$

is a closed, unbounded subset of $[0\,,\,\infty)$ containing 0. What we need for (a) is that there exists a proper ultra-metric d' on X that generates the same topology as d and such that the closure of the value set $\Lambda_{d'}$ of that metric coincides with D. This metric property is proved in the lecture notes by BENDIKOV [8, §2]. Given μ, the Laplacian associated with the standard (d', μ)-processes has the required property by Corollary 7.4. A proof of the additional statement (b) on the ultra-metric can also be found in [8, §2]. ■

It is known that any continuous symmetric Markov semigroup can be extended to all spaces L^p, $1 \leq p < \infty$, as a continuous contraction semigroup. In particular, this is true for the semigroup (P^t). We use the same notation for the extended semigroup and denote by $(-\mathcal{L}_p, \mathsf{Dom}_{\mathcal{L}_p})$ its infinitesimal generator.

Theorem 7.8 *For any* $1 \leq p < \infty$,

$$\mathrm{Spec}(\mathcal{L}_p, \mathsf{Dom}_{\mathcal{L}_p}) = \mathrm{Spec}(\mathcal{L}_2, \mathsf{Dom}_{\mathcal{L}_2}).$$

Proof. Let $\Pi_B : L^2 \to \mathcal{H}_B$ be the orthoprojector onto the space defined in (7.7). Consider the following family of orthoprojectors

$$E_s = \sum_{B \in \mathcal{B} \, : \, r^*(B) > 1/s} \Pi_B .$$

An immediate consequence of (7.9) is that for any $f \in \mathsf{Dom}_{\mathcal{L}_2}$,

$$(\mathcal{L}_2 f, f) = \int_0^\infty s \, d(E_s f, f). \tag{7.11}$$

Claim. For any $f \in L^2$,

$$E_s f = \mathrm{Q}_{1/s} f. \tag{7.12}$$

Indeed, the linear subspace $\mathrm{Q}_{1/s}(L^2) \subset L^2$ consists of all functions in L^2 each of which is constant on any ball B of radius $1/s$. On the other hand, the linear subspace $E_s(L^2) \subset L^2$ can be represented as

$$E_s(L^2) = \sum_{B \in \mathcal{B} \, : \, r^*(B) > 1/s} \Pi_B(L^2) = \sum_{r^*(B) > 1/s} \mathcal{H}_B,$$

It follows that $E_s(L^2)$ is spanned by the set $\{f_B : r^*(B) > 1/s\}$. All those functions take constant values on any ball B of radius $1/s$. Thus $E_s(L^2) \subset \mathrm{Q}_{1/s}(L^2)$. For any ball B of radius $r^*(B) > 1/s$ we can write

$$\mathbf{1}_B = \mu(B) \sum_{C \in \mathcal{B} \, : \, B \subseteq C} f_C \, .$$

It follows that $\mathrm{Q}_{1/s}(\mathcal{D}) \subset E_s(L^2)$, whence $\mathrm{Q}_{1/s}(L^2) \subset E_s(L^2)$ as well. Thus $E_s(L^2) = \mathrm{Q}_{1/s}(L^2)$. Since both E_s and $\mathrm{Q}_{1/s}$ are orthoprojectors, they coincide. This proves the claim.

Since each eigenfunction f_B is compactly supported,

$$\Sigma := \mathrm{Spec}(\mathcal{L}_2, \mathsf{Dom}_{\mathcal{L}_2}) \subset \mathrm{Spec}(\mathcal{L}_p, \mathsf{Dom}_{\mathcal{L}_p}).$$

Let $\lambda \notin \Sigma$, so that $\delta = \min\{|\lambda - s| : s \in \Sigma\} > 0$. Let f and g be bounded functions supported by some ball $B = B_\rho(a)$, where $a \in X$ and $\rho \in \Lambda_*(a)$. Using (7.11), we can write

$$\begin{aligned} \big((\mathcal{L}_2 - \lambda \cdot \mathrm{id})^{-1} f, g\big) &= \int_{[0\,,\,\infty)} \frac{d(E_s f, g)}{s - \lambda} \\ &= \left(\int_{[0\,,\,1/\rho)} + \int_{[1/\rho\,,\,\infty)} \right) \frac{d(E_s f, g)}{s - \lambda} = I_1 + I_2 \, . \end{aligned}$$

(I) *Estimate of* I_1 .

For $s < 1/\rho$ we have evidently $B = B_\rho(a) \subset B_{1/s}(a)$, whence

$$(E_s f, g) = \frac{1}{V(a, 1/s)} (f, 1)(g, 1) = N(a, s) \, (f, 1)(g, 1).$$

It follows that

$$I_1 = (f, 1)(g, 1) \int_{[0\,,\,1/\rho)} \frac{1}{s - \lambda} \, dN(a, s).$$

We use this equality for conjugate exponents (p,q) and obtain

$$|I_1| \le \|f\|_{L^p}\, V(a,\rho)^{\frac{1}{p}}\ \|g\|_{L^q}\, V(a,\rho)^{\frac{1}{q}} \frac{\delta^{-1}}{V(a,1/\tau)} = \delta^{-1}\, \|f\|_{L^p}\, \|g\|_{L^q}\,.$$

(II) *Estimate of* I_2 *.*

We observe that by (7.12)), the signed measure

$$d\mathcal{E}(s) = \mathbf{1}_{[1/r\,,\infty)} d(E_s f, g)$$

is concentrated on the set

$$M_\rho = \left\{ \frac{1}{r} : r \in \Lambda_*(a)\,,\ r \le \rho \right\}.$$

Once more, write $\Lambda_*(a) = \{0\} \cup \{r_k\}$, where k varies as in (2.2), but the increasing sequence $\{r_k\}$ refers to d_*. By assumption, there is i such that $\rho = r_i$, and if we re-parametrize $s_k = 1/r_{i-k}$ then

$$M_\rho = \left\{ \frac{1}{r_k} : k \le i \right\} = \{s_k : k \ge 0\}\,.$$

When X is discrete, M_ρ is finite. Otherwise, $s_k \to \infty$ as $k \to \infty$.

Having these observations in mind, we write

$$I_2 = \sum_{k \ge 0} \frac{1}{s_k - \lambda} \Big((E_{s_{k+1}} f, g) - (E_{s_k} f, g) \Big).$$

Next we apply the Abel transformation with $s_{-1} = 0$ and obtain

$$I_2 = \frac{1}{\lambda}(E_{1/\rho} f, g) + \sum_{k \ge 0} \left(\frac{1}{s_{k-1} - \lambda} - \frac{1}{s_k - \lambda} \right) (E_{s_k} f, g).$$

Since $\lambda \notin \Sigma$, we do not have $\lambda = s_k$ for any k. Thus, there is $k_0 \ge 0$ such that

$$s_{k_0 - 1} < \lambda < s_{k_0}\,,$$

and we write

$$\begin{aligned} I_2 \ &= \ \frac{1}{\lambda}(E_{1/\rho} f, g) + \sum_{k \le k_0} \left(\frac{1}{s_{k-1} - \lambda} - \frac{1}{s_k - \lambda} \right) (E_{s_k} f, g) + \\ &\quad + \sum_{k > k_0} \left(\frac{1}{s_{k-1} - \lambda} - \frac{1}{s_k - \lambda} \right) (E_{s_k} f, g). \end{aligned}$$

By (7.12), $E_s = P_{B_{1/\lambda}}$ is a Markov operator, whence

$$|(E_s f, g)| \le \|f\|_{L^p}\, \|g\|_{L^q}\,.$$

We note that for any $k \ge k_0 + 1$ we have $\frac{1}{s_{k-1} - \lambda} - \frac{1}{s_k - \lambda} > 0$. After all these preparations we can estimate I_2 :

$$\begin{aligned} |I_2| \ &\le \ \left(\frac{1}{\lambda} + 2(k_0+1)\delta^{-1} \right) \|f\|_{L^p}\, \|g\|_{L^q} + \sum_{k > k_0} \left(\frac{1}{s_{k-1} - \lambda} - \frac{1}{s_k - \lambda} \right) \big|(E_{s_k} f, g)\big| \\ &\le \ \left(\frac{1}{\lambda} + 2(k_0+1)\delta^{-1} + \frac{1}{s_{k_0} - \lambda} \right) \|f\|_{L^p}\, \|g\|_{L^q} \\ &\le \ \left(\frac{1}{\lambda} + (2k_0+3)\delta^{-1} \right) \|f\|_{L^p}\, \|g\|_{L^q}\,. \end{aligned}$$

Putting together the bounds obtained in (I) and (II), we finally come to the desired inequality

$$\left|\Big((\mathcal{L}_2-\lambda\ \mathrm{id})^{-1}f,g\Big)\right| \le \left(\frac{1}{\lambda}+(2k_0+4)\delta^{-1}\right)\|f\|_{L^p}\,\|g\|_{L^q}\,.$$

This inequality shows that the operator $(\mathcal{L}_2-\lambda\ \mathrm{id})^{-1}$ can be extended to L^p as a bounded operator. The extended operator evidently coincides with $(\mathcal{L}_p-\lambda\,\mathrm{id})^{-1}$. Thus $\lambda\notin \mathrm{Spec}(\mathcal{L}_p,\mathsf{Dom}_{\mathcal{L}_p})$ and therefore

$$\mathrm{Spec}(\mathcal{L}_p,\mathsf{Dom}_{\mathcal{L}_p})\subseteq \mathrm{Spec}(\mathcal{L}_2,\mathsf{Dom}_{\mathcal{L}_2}).$$

The proof of the theorem is finished. ■

We remark that (7.11) and (7.12) indeed provide explicitly the spectral resolution, as outlined in the Introduction.

The last theorem of this section concerns harmonic functions. Notice that the semigroup $(P^t)_{t>0}$ admits an extension to L^∞ as a contraction semigroup, but this extension is not a continuous semigroup unless X is discrete. We define the Laplacian $(\mathcal{L}_\infty,\mathsf{Dom}_{\mathcal{L}_\infty})$ as a weak infinitesimal generator, that is, we define $\mathsf{Dom}_{\mathcal{L}_\infty}\subset L^\infty$ as the set of functions f such that

$$\mathcal{L}_\infty f(x)=\lim_{t\to 0}\frac{f(x)-P^tf(x)}{t}$$

exists for all $x\in X$ and belongs to L^∞. This defines $\mathcal{L}_\infty f(x)$ for $f\in\mathsf{Dom}_{\mathcal{L}_\infty}$ and $x\in X$.

Theorem 7.9 (Strong Liouville property) *Any measurable function $f:X\to[0\,,\infty]$ which satisfies $Pf=f$ must be constant.*

In particular, $0\in\mathrm{Spec}(\mathcal{L}_\infty,\mathsf{Dom}_{\mathcal{L}_\infty})$ is an eigenvalue of multiplicity 1.

Proof. We may assume without loss of generality that we are dealing with the standard (d_*,μ)-process on X, where $d=d_*$ is the intrinsic metric and the distance distribution is σ_*, the inverse exponential distribution (1.5) .

Step 1. We first prove the claim assuming that f takes only finite values.

By assumption,

$$f=\int_0^\infty \mathrm{Q}_s f\,d\sigma_*(s), \tag{7.13}$$

see (1.3). For any fixed $r>0$ we apply to both sides of this equation the operator Q_r. Using (2.6), we obtain

$$\mathrm{Q}_rf=\int_0^\infty \mathrm{Q}_{\max\{r,s\}}f\,d\sigma_*(s)=\sigma_*(r)\mathrm{Q}_rf+\int_r^\infty \mathrm{Q}_sf\,d\sigma_*(s),$$

whence,

$$\big(1-\sigma_*(r)\big)\mathrm{Q}_rf=\int_r^\infty \mathrm{Q}_sf\,d\sigma_*(s). \tag{7.14}$$

Let $B\in\mathcal{B}$ be a ball with diameter r, and let r' be the diameter of the predecessor ball B'. We consider (7.14) for r and for r', and take the difference: for any $x\in B$,

$$\big(1-\sigma_*(r)\big)\mathrm{Q}_rf(x)-\big(1-\sigma_*(r')\big)\mathrm{Q}_{r'}f(x)=\int_r^{r'}\mathrm{Q}_sf(x)d\sigma_*(s)=\big(\sigma_*(r')-\sigma_*(r)\big)\mathrm{Q}_rf(x),$$

or equivalently

$$\big(1-\sigma_*(r')\big)\,\big(\mathrm{Q}_rf(x)-\mathrm{Q}_{r'}f(x)\big)=0.$$

We conclude that

$$\mathrm{Q}_rf(x)=\mathrm{Q}_{r'}f(x) \tag{7.15}$$

for every $x \in B$ and all $r, r' \in \Lambda_*(x)$, where initially r' is the smallest element in $\Lambda_*(x)$ with $r' > r$. Inductively, we get that (7.15) holds for all $r, r' \in \Lambda_*(x)$ without further restriction, that is, the function $r \mapsto \mathrm{Q}_r f(x)$ is constant. Since $x \mapsto \mathrm{Q}_r f(x)$ is constant on each ball of radius r, we get that the latter function is constant both in x and in r. Since $f(x) = Pf(x)$, f is constant as well.

Step 2. Assume now that $f(a) < \infty$ for some $a \in X$. Let $B = B_r(a)$ and $B' = B_{r'}(a)$ be as above in Step 1. By (7.13),

$$\big(\sigma_*(r') - \sigma_*(r)\big) P_r f(a) \int_r^{r'} \mathrm{Q}_s f(a)\, d\sigma_*(s) \le \int_0^\infty \mathrm{Q}_s f(a)\, d\sigma_*(s) = f(a) < \infty\,.$$

Thus, $\mathrm{Q}_r f(a) < \infty$. Recalling the structure (2.2) of $\Lambda_*(a)$, again by induction, $\mathrm{Q}_r f(a) < \infty$ for any $r > 0$. In particular, for any $r > 0$, the function $f_r(x) = \mathrm{Q}_r f(x)$ takes only finite values and evidently satisfies the equation $Pf_r = f_r$. By the first part of the proof, f_r is constant. The identity (7.13) now yields that f is constant.

For the last statement of the theorem, assume that $f \in \mathsf{Dom}_{\mathcal{L}_\infty}$ and $\mathcal{L}_\infty f = 0$. Since f is a bounded function we can assume that it is non-negative. Since $f \in \mathsf{Dom}_{\mathcal{L}_\infty}$ we can write

$$Pf - f = -\int_0^1 \mathcal{L}_\infty \mathrm{Q}_s f\, ds = -\int_0^1 \mathrm{Q}_s \mathcal{L}_\infty f\, ds = 0,$$

whence, by Step 1, f is constant. This concludes the proof. ■

Corollary 7.10 *Assume that for some measure μ' the semigroup $\{P^t\}_{t>0}$ is symmetric in $L^2(\mu')$. Then μ' is proportional to μ.*

8 The p-adic fractional derivative

Consider the field $\mathbb{Q}_p$ of p-adic numbers endowed with the p-adic norm $\|x\|_p$ and the p-adic ultra-metric $d(x,y) = d_p(x,y) = \|x-y\|_p$. Let μ_p be the Haar measure on $\mathbb{Q}_p$, normalized such that $\mu_p(\mathbb{Z}_p) = 1$. Let $\mathcal{D}$ be the set of test functions on $\mathbb{Q}_p$, that is, locally constant functions with compact support.

The notion of p-adic fractional derivative, closely related to the concept of p-adic Quantum Mechanics, was introduced in several mathematical papers by VLADIMIROV [42], VLADIMIROV AND VOLOVICH [43] and VLADIMIROV, VOLOVICH AND ZELENOV [44]. In particular, in [42] a one-parametric family $\{(D^\alpha, \mathcal{D})\}_{\alpha>0}$ of operators, called operators of fractional derivative of order α, has been introduced.

Definition 8.1 The operator $(D^\alpha, \mathcal{D})$, $\alpha > 0$, is defined via the Fourier transform on the locally compact Abelian group $\mathbb{Q}_p$ by

$$\widehat{D^\alpha f}(\xi) = \|\xi\|_p^\alpha\, \widehat{f}(\xi),\ \ \xi \in \mathbb{Q}_p. \tag{8.1}$$

It was shown by the above authors that each operator $(D^\alpha, \mathcal{D})$ can be written as a Riemann-Liouville type singular integral operator

$$D^\alpha f(x) = \frac{p^\alpha - 1}{1 - p^{-\alpha-1}} \int_{\mathbb{Q}_p} \frac{f(x) - f(y)}{\|x-y\|_p^{1+\alpha}}\, d\mu_p(y). \tag{8.2}$$

The aim of this section is to show that the operator $(D^\alpha, \mathcal{D})$ is in fact restricted to $\mathcal{D}$ Laplacian (minus the Markov generator) of an appropriate isotropic Markov semigroup $\{P_\alpha^t\}_{t>0}$, as constructed and studied throughout this paper. We use the step length distribution $\sigma_\alpha = \sigma_{\alpha,p}$ with distribution function

$$\sigma_\alpha(r) = \begin{cases} \exp\left(-\left(\frac{p}{r}\right)^\alpha\right) & \text{if} \quad r > 0\,, \\ 0 & \text{if} \quad s \le 0\,. \end{cases}$$

We now consider the $(d_p, \mu_p, \sigma_\alpha)$-process and its Markov semigroup $(P_\alpha^t)_{t>0}$, as constructed in (1.2) – (1.4),

$$P_\alpha^t f(x) = \int_0^\infty \mathrm{Q}_r f(x)\, d\sigma_\alpha^t(r).$$

The semigroup acts in L^2, and we let $(\mathcal{L}^\alpha\,, \mathsf{Dom}_{\mathcal{L}^\alpha})$ be the corresponding Laplacian.

Theorem 8.2 *For any* $\alpha > 0$,

$$(\mathcal{L}^\alpha\,, \mathcal{D}) = (D^\alpha\,, \mathcal{D}).$$

Proof. By Theorem (7.2), for $f \in \mathcal{D}$,

$$\mathcal{L}^\alpha f(x) = \int_{\mathbb{Q}_p} \big(f(x) - f(y)\big) J_\alpha(x,y) d\mu_p(y),$$

whence we are left to compute the function

$$J_\alpha(x,y) = \int_0^{1/d_*(x,y)} N(x,\tau)\, d\tau = \int_0^{1/d_*(x,y)} \frac{d\tau}{\mu_p\big(B^*_{1/\tau}(x)\big)},$$

where $B^*_{1/\tau}(x)$ is the d_*- ball at x of radius $1/\tau$. By Example (4.11),

$$d_*(x,y) = \left(\frac{\|x-y\|_p}{p}\right)^\alpha,$$

whence

$$B^*_{1/\tau}(x) = B_{p/\tau^{1/\alpha}}(x),$$

where $B_{p/\tau^{1/\alpha}}(x)$ is the d_p-ball at x of radius $p/\tau^{1/\alpha}$. Putting all these facts together we write

$$J_\alpha(x,y) = \int_0^{\left(p/\|x-y\|_p\right)^\alpha} \frac{d\tau}{\mu_p\big(B_{p/\tau^{1/\alpha}}(x)\big)} = p^\alpha \int_{\|x-y\|_p}^\infty \frac{\alpha r^{-\alpha-1}\, dr}{\mu_p\big(B_r(x)\big)}.$$

The value set of the metric is $\Lambda_{d_p}(x) = \{p^k : k \in \mathbb{Z}\}$ for every $x \in \mathbb{Q}_p$. Thus, if $\|x-y\|_p = p^k$ then the last integral is

$$\begin{aligned} \int_{p^k}^\infty \frac{\alpha r^{-\alpha-1}\, dr}{\mu\big(B_r(x)\big)} &= \sum_{n\ge k} \int_{p^n}^{p^{n+1}} \frac{\alpha r^{-\alpha-1}\, dr}{\mu\big(B_r(x)\big)} \\ &= \sum_{n\ge k} \int_{p^n}^{p^{n+1}} \frac{\alpha r^{-\alpha-1}\, dr}{p^n} = \sum_{n\ge k} \frac{1}{p^n}\left(\frac{1}{p^{n\alpha}} - \frac{1}{p^{(n+1)\alpha}}\right) \\ &= \left(1 - \frac{1}{p^\alpha}\right) \sum_{n\ge k} \frac{1}{p^{n(\alpha+1)}} = \left(1 - \frac{1}{p^\alpha}\right) \frac{p^{-k(\alpha+1)}}{1-p^{-(\alpha+1)}} \\ &= \frac{1-p^{-\alpha}}{1-p^{-(\alpha+1)}} \left(\frac{1}{p^k}\right)^{\alpha+1} = \frac{1-p^{-\alpha}}{1-p^{-(\alpha+1)}} \left(\frac{1}{\|x-y\|_p}\right)^{\alpha+1}. \end{aligned}$$

Thus finally we obtain the following equality

$$J_\alpha(x,y) = \frac{p^\alpha - 1}{1 - p^{-(\alpha+1)}} \left(\frac{1}{\|x-y\|_p} \right)^{\alpha+1}. \tag{8.3}$$

In view of (8.2), this proves the claim. ■

As a consequence of Theorem 8.2, Corollary 4.5 and Example 4.11 we obtain that the following heat kernel estimate hold.

Theorem 8.3 *The semigroup* $(P^t_\alpha)_{t>0}$ *admits a continuous transition density* $p_\alpha(t,x,y)$ *with respect to Haar measure* μ_p *which satisfies*

$$\frac{c_1\, t}{(t^{1/\alpha} + \|x-y\|_p)^{1+\alpha}} \le p_\alpha(t,x,y) \le \frac{c_2\, t}{(t^{1/\alpha} + \|x-y\|_p)^{1+\alpha}},$$

for some constants $c_1\,, c_2 > 0$.

Theorem 8.4 *The semigroup* $(P^t_\alpha)_{t>0}$ *is transient if and only if* $0 < \alpha < 1$. *In the transient case, its Green function is given explicitly as*

$$G^\alpha(x,y) = \frac{1-p^{-\alpha}}{1-p^{\alpha-1}} \left(\frac{1}{\|x-y\|_p} \right)^{-\alpha+1}.$$

Proof. The characterization of transience follows from Example 5.7. In this case, by Theorem 5.4(3)

$$\begin{aligned} G^\alpha(x,y) &= \int_0^{1/d_*(x,y)} \frac{N(x,\tau)\,d\tau}{\tau^2} = \int_{d_*(x,y)}^\infty N\left(x, \frac{1}{\tau}\right) d\tau \\ &= \int_{d_*(x,y)}^\infty \frac{d\tau}{V(x,\tau)} = \int_{(\|x-y\|_p/p)^\alpha}^\infty \frac{d\tau}{\mu\big(B_{p\tau^{1/\alpha}}(x)\big)}. \end{aligned}$$

With the change of variables $s = p\tau^{1/\alpha}$, if $\|x-y\|_p = p^k$ for $k \in \mathbb{Z}$, we obtain

$$\begin{aligned} G^\alpha(x,y) &= \frac{1}{p^\alpha} \int_{p^k}^\infty \frac{\alpha s^{\alpha-1}\,ds}{\mu(B_s(x))} = \frac{1}{p^\alpha} \sum_{n\ge k} \int_{p^n}^{p^{n+1}} \frac{\alpha s^{\alpha-1}\,ds}{p^n} \\ &= \frac{1}{p^\alpha} \sum_{n \ge k} \frac{1}{p^n} \left(p^{(n+1)\alpha} - p^{n\alpha} \right) = \frac{p^\alpha - 1}{p^\alpha(1-p^{\alpha-1})} \left(\frac{1}{p^k} \right)^{1-\alpha} \\ &= \frac{1-p^{-\alpha}}{1-p^{\alpha-1}} \left(\frac{1}{\|x-y\|_p} \right)^{-\alpha+1}. \end{aligned}$$

The proof is finished. ■

Let $\{X_t\}$ be the Markov process on $\mathbb{Q}_p$ driven by the Markov semigroup $(P^t_\alpha)_{t>0}$. The semigroup $\{P^t_\alpha\}_{t>0}$ is translation invariant, whence the process has independent and stationary increments. For any given $\gamma > 0$ and $t > 0$, consider the moment of order γ of $\mathcal{X}_t$ defined in terms of the p-adic distance $d_p(x,y)$:

$$\mathcal{M}_\gamma(t) = \mathbb{E}(\|\mathcal{X}_t\|_p^\gamma),$$

where as in the Section 6, $\mathbb{E}$ is expectation with respect to the probability measure on the trajectory space of the process starting at 0. Applying Theorem 6.3 we obtain the following estimates.

Theorem 8.5 *The moment $\mathcal{M}_\gamma(t)$ is finite if and only if $\gamma < \alpha$. Moreover, if $\gamma < \alpha$, there exists a constant $\kappa = \kappa(\alpha) > 0$ such that*

$$\frac{\kappa t^{\gamma/\alpha}}{\alpha - \gamma} \le \mathcal{M}_\gamma(t) \le \frac{\alpha t^{\gamma/\alpha}}{\alpha - \gamma}.$$

Let D_q^α be the minus infinitesimal generator of the semigroup $(P_\alpha^t)_{t>0}$ acting in $L^q(\mu_p)$, $1 \le q < \infty$. Applying Theorems 7.3 and 7.8) we obtain the following spectral result.

Theorem 8.6 *For any $\alpha > 0$ and $1 \le q < \infty$,*

$$\mathrm{Spec}(D_q^\alpha) = \{p^{k\alpha} : k \in \mathbb{Z}\} \cup \{0\}.$$

Each $\lambda_k = p^{k\alpha}$ is an eigenvalue having infinite multiplicity.

In the general setting of Theorems 7.3 and 7.8, some eigenvalues may well have finite multiplicity and some not. Indeed, attached to each ball B of d_*-diameter $1/\lambda$ there are the eigenvalue λ and the corresponding finite dimensional eigenspace $\mathcal{H}_B$. This eigenspace is spanned by the finitely many functions

$$f_C = \frac{1}{\mu(C)}\mathbf{1}_C - \frac{1}{\mu(B)}\mathbf{1}_B\,,$$

where C runs through all balls whose predecessor (in the sense of Definition 2.1) is $C' = B$. Recall that $\dim \mathcal{H}_B = l(B) - 1$, where $l(B) = \sharp\{C \in \mathcal{B} : C' = B\}$.

It follows that in general, if there exists only finite number of distinct balls of d_*-diameter $1/\lambda$ then the eigenvalue λ has finite multiplicity.

This is clearly not the case of the ultra-metric measure space $(\mathbb{Q}_p, d_p, \mu_p)$ and the semigroup $(P_\alpha^t)_{t>0}$. Indeed, every d_*-ball has its diameter in the set $\Lambda_\alpha = \{p^{k\alpha} : k \in \mathbb{Z}\}$, and each ball $B_r(0)$ has infinitely many disjoint translates $\{a_i + B_r(0)\}_{i=1}^\infty$, which cover $\mathbb{Q}_p$ and are balls of the same diameter. Thus, all eigenvalues have infinite multiplicity.

Remark 8.7 Let $\mathcal{H}(\lambda)$ be the eigenspace corresponding to the eigenvalue $\lambda \in \Lambda_\alpha$. Then

$$L^2 = \bigoplus_{\lambda \in \Lambda_\alpha} \mathcal{H}(\lambda) \quad \text{and} \quad \mathcal{H}(\lambda) = \bigoplus_{i=1}^{\infty} \mathcal{H}_{a_i + B_{1/\lambda}(0)}. \tag{8.4}$$

As after (7.8), we choose for each closed ball $B \subset \mathbb{Q}_p$ an orthonormal basis $\{e_i^B : 1 \le i \le p-1\}$ in $\mathcal{H}_B$. In view of (8.4), the set of eigenfunctions $\{e_i^B : B \in \mathcal{B}, 1 \le i \le p-1\}$ is an orthonormal basis in L^2. (This reasoning applies to arbitrary ultrametric spaces.) Whether this set is a Schauder basis in L^q, $1 \le q < \infty$, is an open question.

Our next result (see Theorem 8.8 below) can be viewed as a p-adic version of the Polya's theorem: a function $F(|\xi|), \xi \in \mathbb{R}$, is the characteristic function of a symmetric probability density function whenever $F(0) = 1$, $F(s)$ is convex for $s > 0$ and $F(+\infty) = 0$.

Let $(P^t)_{t>0}$ be isotropic Markov semigroup on the ultra-metric measure space $(\mathbb{Q}_p, d_p, \mu_p)$ as constructed in (1.2) – (1.4). This semigroup acts in C_0 - the Banach space of continuous functions vanishing at ∞. Moreover this semigroup is translation invariant. It follows that there exists a weakly continuous convolution semigroup $(p_t)_{t>0}$ of probability measures on $\mathbb{Q}_p$ such that

$$P^t f(x) = p_t * f(x). \tag{8.5}$$

As the semigroup $(P^t)_{t>0}$ admits a continuous transition density $p(t,x,y)$ with respect to the Haar measure μ_p, all the measures p_t are absolutely continuous with respect to the measure μ_p and have continuous densities $p_t(y) = p(t,0,y)$ given by

$$p_t(y) = \int_0^\infty q_s(y) d\sigma(s)^t, \text{ where } q_s(y) = \frac{1_{B_s(0)}(y)}{\mu_p(B_s(0))}.$$

To find the Fourier transform $\widehat{p_t}(\xi)$ of the measure p_t we argue as follows. The ball $B_s(0)$, $p^k \leq s < p^{k+1}$, is a compact subgroup $p^{-k}\mathbb{Z}_p$ of the group $\mathbb{Q}_p$, whence the measure $\omega_s = q_s\mu_p$ coincides with the normed Haar measure of the compact subgroup $p^{-k}\mathbb{Z}_p$. As in the general setting of locally compact Abelian groups the Fourier transform of the normed Haar measure of any compact subgroup is the indicator of its annihilator group and, in our particular case, the annihilator of the group $p^{-k}\mathbb{Z}_p$ is the group $p^k\mathbb{Z}_p$ we obtain

$$\widehat{\omega_s}(\xi) = 1_{p^k\mathbb{Z}_p}(\xi) = 1_{[0,p^{-k}]}(\|\xi\|_p), \text{ where } p^k \leq s < p^{k+1}.$$

It follows that, when $\|\xi\|_p = p^{-l}, l \in \mathbb{Z}$,

$$\begin{aligned} \widehat{p_t}(\xi) &= \sum_{k:k\leq l} \left(\sigma(p^{k+1})^t - \sigma(p^k)^t\right) \\ &= \sigma(p^{l+1})^t = \sigma\left(\frac{p}{\|\xi\|_p}\right)^t. \end{aligned}$$

Choosing a decreasing bijection $F : [0,\infty) \to (0,1]$ which coincides with the function $\tau \to \sigma(p/\tau)$ at points $\tau = p^l, l \in \mathbb{Z}$, we finally write

$$\widehat{p_t}(\xi) = F(\|\xi\|_p)^t. \tag{8.6}$$

Conversely, if a decreasing bijection $F : [0,\infty) \to (0,1]$ is given, writing $F(s) = \exp(-\psi(s))$ and applying the results of Theorem 8.2 (the case $\alpha = 1$) and Theorem 7.5 we conclude that there exists a weakly continuous convolution semigroup $(p_t^F)_{t>0}$ of probability measures on $\mathbb{Q}_p$ and an isotropic Markov semigroup $(P_F^t)_{t>0}$ as constructed in (1.2) – (1.4) such that

$$\widehat{p_t^F}(\xi) = F(\|\xi\|_p)^t \tag{8.7}$$

and

$$P_F^t f(x) = p_t^F * f(x). \tag{8.8}$$

Thus finally we obtain the following result.

Theorem 8.8 *The equations (8.5) and (8.6) when F runs over all decreasing bijections* $[0,\infty) \to (0,1]$ *give a complete description of the class of isotropic Markov semigroups on the ultra-metric measure space* $(\mathbb{Q}_p, d_p, \mu_p)$ *as constructed in (1.2) – (1.4).*

Recall for comparison that defined on the Abelian group $\mathbb{R}^d$ (equipped with the Euclidean metric) the function $F(\|\xi\|) = \exp(-\|\xi\|^\alpha)$, $\alpha > 0$, can be represented as the Fourier transform of a probability measure defined on $\mathbb{R}^d$ if and only if $\alpha \leq 2$.

9 Vladimirov's Laplacian

We now consider the linear space $\mathbb{Q}_p^n = \mathbb{Q}_p \times ... \times \mathbb{Q}_p$ (n times) over the field of p-adic numbers, equipped with the ultra-metric

$$d_{p,\alpha}(x,y) = \|x-y\|_{p,\alpha} ,$$

where $\alpha = (\alpha_1, \alpha_2, ..., \alpha_n)$ is any given n-tuple with entries $\alpha_i > 0$ and

$$\|z\|_{p,\alpha} = \max\left\{ \|z_i\|_p^{\alpha_i} : i = 1, 2, ..., n \right\}. \tag{9.1}$$

We set $\|z\|_p = \max\left\{ \|z_i\|_p : i = 1, 2, ..., n \right\}$ and observe that $\|z\|_p$ is a norm, that is, $\|az\|_p = \|a\|_p \|z\|_p$.

For any given n-tuple $\alpha = (\alpha_1, \ldots, \alpha_n)$ set $\alpha_* = \max\{\alpha_i\}$ and observe that the identity map

$$i : \left(\mathbb{Q}_p^n, (d_{p,\alpha})^{1/\alpha_*} \right) \to \left(\mathbb{Q}_p^n, \|\cdot\|_p \right)$$

is a homeomorphism but not a quasi-isometry unless all entries $\alpha_i = \beta$. This fact plays an essential role in the study of the class of Laplacians introduced below.

Definition 9.1 On the set $\mathcal{D}$ of test functions on $\mathbb{Q}_p^n$ we define the operator

$$D^\alpha f(x) = \sum_{i=1}^n D_{x_i}^{\alpha_i} f(x), \quad \text{where} \quad x = (x_1, x_2, ..., x_n). \tag{9.2}$$

(The index x_i refers to the action on the i-th variable.)

An operator of this type, namely the operator D^α in $\mathbb{Q}_p^3$ with $\alpha = (2,2,2)$, was introduced by Vladimirov [42] as an analogue of $-\Delta$, where Δ is the classical Laplace operator in $\mathbb{R}^3$. This operator, which we denote $\mathfrak{D}^2$ is translation invariant and homogeneous, that is,

$$\mathfrak{D}^2 \tau_y(f) = \tau_y(\mathfrak{D}^2 f), \quad \text{where} \quad \tau_y f(x) = f(x+y).$$

and

$$\mathfrak{D}^2 \theta_a(f) = \|a\|_p^2 \theta_a(\mathfrak{D}^2 f), \quad \text{where} \quad \theta_a f(x) = f(ax_1, ax_2, ax_3).$$

It follows that the Green function $G(x,y)$ of the operator $\mathfrak{D}^2$ is also translation invariant and homogeneous:

$$G(x,y) = G(x-z, y-z) \quad \text{and} \quad G(ax, ay) = G(x,y)/\|a\|_p .$$

In particular, setting $\mathfrak{E}(x) = G(x,0)$, we will obtain the following identity

$$\mathfrak{E}(a,a,a) = \frac{\mathfrak{E}(1,1,1)}{\|a\|_p}.$$

This identity was observed in [42]. It gives an idea of how the Green function of the operator $\mathfrak{D}^2$ (in Vladimirov's terminology, the fundamental solution of the equation $\mathfrak{D}^2\mathfrak{E} = \delta$) behaves at infinity/at zero.

Below in Proposition 9.4, we will prove that

$$\mathfrak{E}(a_1, a_2, a_3) \simeq \frac{1}{\|a\|_p}, \tag{9.3}$$

where $a = (a_1, a_2, a_3)$ and $\|a\|_p = \max\{\|a_i\|_p\}$.

We shall extend the asymptotic property (9.3) to the more general operators D^α. In general the homogeneity property is lost, whence we will develop some tools as a compensation.

Let $\mu_p = \bigotimes \mu_{p,i}$ be the additive Haar measure on $\mathbb{Q}_p^n$ and let $L^2 = L^2(\mathbb{Q}_p^n, \mu_p)$. We list some properties of the operator $(D^\alpha, \mathcal{D})$ of (9.2) which follow directly from the corresponding properties of the "one-dimensional" Laplacians D^{α_i}.

1. $(D^\alpha, \mathcal{D})$ is a non-negative definite symmetric operator.
2. $(D^\alpha, \mathcal{D})$ admits a complete system of compactly supported eigenfunctions. In particular, the operator $(D^\alpha, \mathcal{D})$ is essentially self-adjoint.
3. The semigroup $P_\alpha^t = \exp(-tD^\alpha)$ is symmetric and Markovian. It admits a transition density (heat kernel) $p_\alpha(t,x,y)$ which has the following form
$$p_\alpha(t,x,y) = \prod_{i=1}^n p_{\alpha_i}(t,x_i,y_i).$$
In particular, for all $x \in \mathbb{Q}_p^n$ and all $t>0$,
$$c_1\, t^{-A} \le p_\alpha(t,x,x) \le c_2\, t^{-A},$$
where
$$A = \sum_{i=1}^n \frac{1}{\alpha_i}$$
and $c_1, c_2 > 0$ are some constants.
4. The semigroup $\{P_\alpha^t\}_{t>0}$ is transient if and only if $A > 1$.

Our next goal is to find off-diagonal estimates of the heat kernel $p_\alpha(t,x,y)$ and, in the transient case, the Green function $G^\alpha(x,y)$ in metric terms. Observe that thanks to the group structure of $\mathbb{Q}_p^n$, the functions $(x,y) \mapsto p_\alpha(t,x,y)$ and $(x,y) \mapsto G^\alpha(x,y)$ are translation invariant. Hence, setting
$$p_\alpha(t,z) = p_\alpha(t,z,0) \quad \text{and} \quad G^\alpha(z) = G^\alpha(z,0)$$
we obtain
$$p_\alpha(t,x,y) = p_\alpha(t,x-y) \quad \text{and} \quad G^\alpha(x,y) = G^\alpha(x-y).$$

Proposition 9.2 *The heat kernel satisfies*
$$p_\alpha(t,z) \simeq t^{-A} \prod_{i=1}^n \min\left\{1, \frac{t^{1+1/\alpha_i}}{\|z_i\|_p^{1+\alpha_i}}\right\}$$
uniformly for all $t>0$ and $z \in \mathbb{Q}_p^n$.

In particular, there are $c_1, c_2 > 0$ such that for all $t > \|z\|_{p,\alpha}$,
$$c_1\, t^{-A} \le p_\alpha(t,z) \le c_2\, t^{-A} \tag{9.4}$$

Proof. We recall the definition 9.1 of $\|z\|_{p,\alpha}$ and apply Theorem 8.3: the "one-dimensional" heat kernels can be estimated as follows.
$$p_{\alpha_i}(t,z_i) \simeq \frac{t}{\left(t^{1/\alpha_i} + \|z_i\|_p\right)^{1+\alpha_i}} \simeq \frac{1}{t^{1/\alpha_i}} \min\left\{1, \frac{t^{1+1/\alpha_i}}{\|z_i\|_p^{1+\alpha_i}}\right\}.$$

This implies the proposed result. ■

Proposition 9.3 *Assume that the semigroup $\{P^t_\alpha\}_{t>0}$ is transient. Then, for all $z \in \mathbb{Q}_p^n$ and some $C_1 > 0$,*

$$G^\alpha(z) \geq C_1 \left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1}.$$

For any $\kappa > 0$, we define the set

$$\Omega(\kappa) = \left\{ x \in \mathbb{Q}_p^n : \max_i \left\{ \|x_i\|_p^{\alpha_i} \right\} \leq \kappa \min_i \left\{ \|x_i\|_p^{\alpha_i} \right\} \right\}.$$

Then, for all $z \in \Omega(\kappa)$ and some constant $C_2 > 0$ which depends on κ,

$$G^\alpha(z) \leq C_2 \left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1}.$$

Proof. Recall that transience holds precisely when $A > 1$. To prove the lower bound, we use (9.4) and write

$$G^\alpha(z) = \int_0^\infty p_\alpha(t,z)\,dt \geq \int_{\|z\|_{p,\alpha}}^\infty p_\alpha(t,z)\,dt \geq C_1 \int_{\|z\|_{p,\alpha}}^\infty t^{-A}\,dt = c_1 \left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1}.$$

On the other hand we have

$$G^\alpha(z) = \left(\int_0^{\|z\|_{p,\alpha}} + \int_{\|z\|_{p,\alpha}}^\infty \right) p_\alpha(t,z)\,dt = I + II.$$

To estimate the second term II, we use again the inequality (9.4),

$$II \simeq \int_{\|z\|_{p,\alpha}}^\infty t^{-A}\,dt \simeq \left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1}.$$

To estimate the first term we use Proposition 9.2,

$$\begin{aligned} I &\leq c \int_0^{\|z\|_{p,\alpha}} t^{-A} \prod_{i=1}^n \frac{t^{1+1/\alpha_i}}{\|z_i\|_p^{1+\alpha_i}}\,dt \\ &= c \int_0^{\|z\|_{p,\alpha}} \prod_{i=1}^n \frac{1}{\|z_i\|_p^{1+\alpha_i}} t^n\,dt = c' \prod_{i=1}^n \frac{1}{\|z_i\|_p^{1+\alpha_i}} \|z\|_{p,\alpha}^{n+1}. \end{aligned}$$

When $z \in \Omega(\kappa)$, we obtain

$$\begin{aligned} I &\leq c'' \prod_{i=1}^n \frac{1}{\|z_i\|_p^{1+\alpha_i}} \left(\min\{\|z_i\|_p^{\alpha_i}\} \right)^{n+1} \leq c'' \min\{\|z_i\|_p^{\alpha_i}\} \prod_{i=1}^n \frac{1}{\|z_i\|_p} \\ &= c'' \min\{\|z_i\|_p^{\alpha_i}\} \prod_{i=1}^n \frac{1}{(\|z_i\|_p^{\alpha_i})^{1/\alpha_i}}. \end{aligned}$$

Next,

$$\prod_{i=1}^n \frac{1}{(\|z_i\|_p^{\alpha_i})^{1/\alpha_i}} \leq \prod_{i=1}^n \frac{1}{\left(\min\{\|z_j\|_p^{\alpha_j}\}\right)^{1/\alpha_i}} = \left(\frac{1}{\min\{\|z_j\|_p^{\alpha_j}\}}\right)^A,$$

whence

$$I \leq c'' \left(\frac{1}{\min\{\|z_j\|_p^{\alpha_j}\}}\right)^{A-1}.$$

Again using the fact that $z \in \Omega(\kappa)$, we write

$$\left(\frac{1}{\min\{\|z_j\|_p^{\alpha_j}\}}\right)^{A-1} \leq \left(\frac{\kappa}{\max\{\|z_j\|_p^{\alpha_j}\}}\right)^{A-1} = c(\kappa)\left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1}.$$

The obtained upper bounds on the integrals I and II imply the desired upper bound for $G^\alpha(z)$. ■

Proposition 9.4 *Let $\alpha = (\alpha_1, \dots, \alpha_n) = (\beta, \dots, \beta)$ be an n-tuple having all entries equal to β. Assume that $(n-1)/2 < \beta < n$. Then the semigroup $\{P_\alpha^t\}$ is transient and the Green function $G^\alpha(z)$ satisfies*

$$c_1\left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1} \leq G^\alpha(z) \leq c_2\left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1},$$

equivalently,

$$c_1\left(\frac{1}{\|z\|_p}\right)^{n-\beta} \leq G^\alpha(z) \leq c_2\left(\frac{1}{\|z\|_p}\right)^{n-\beta},$$

for all $z \in \mathbb{Q}_p^n$ and some $c_1, c_2 > 0$.

Proof. Transience follows from (9.4) because $A = n/\beta > 1$. Then Proposition 9.3 implies the desired lower bound of the Green function

$$G^\alpha(z) \geq c_1\left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1}.$$

To prove the upper bound, we observe that like $\mathbb{D}^2$, the Laplacian D^α is homogeneous, that is

$$D^\alpha \circ \theta_a = \|a\|_p^\beta \cdot \theta_a \circ D^\alpha,$$

for all $a \in \mathbb{Q}_p$.

This implies that also the Green function $G^\alpha(z)$ is homogeneous, that is

$$G^\alpha(az) = \|a\|_p^{n-\beta}\, G^\alpha(z)$$

for all $a \in \mathbb{Q}_p$ and $z \in \mathbb{Q}_p^n$.

Assume now that $\|z\|_{p,\alpha} = \|z_1\|_p^\beta > 0$. Then

$$\begin{aligned} G^\alpha(z) &= G^\alpha\big(z_1(1, z_2/z_1, ..., z_n/z_1)\big) = \|z_1\|_p^{n-\beta}\, G^\alpha(1, z_2/z_1, ..., z_n/z_1) \\ &= \left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1} G^\alpha(1, z_2/z_1, ..., z_n/z_1) \\ &\leq \left(\frac{1}{\|z\|_{p,\alpha}}\right)^{A-1} \sup\big\{G^\alpha(1, x_2, ..., x_n) : x_i \in \mathbb{Z}_p\big\}. \end{aligned}$$

Next we apply our assumption $\beta > (n-1)/2$ and the heat kernel upper bound resulting from Proposition 9.2

$$\begin{aligned} G^\alpha(1, x_2, ..., x_n) &= \int_0^\infty p_\alpha(t, (1, x_2, ..., x_n)\, dt \\ &= \left(\int_0^1 + \int_1^\infty\right) p_\alpha(t, (1, x_2, ..., x_n)\, dt \\ &\leq c\int_0^1 t^{-\frac{n}{\beta}}\, t^{1+\frac{1}{\beta}}\, dt + c'\int_1^\infty t^{-\frac{n}{\beta}}\, dt = c_2 < \infty. \end{aligned}$$

The desired upper bound follows. ■

10 Random walks on a tree and jump processes on its boundary

Rooted trees and their boundaries. A tree is a connected graph T without cycles (closed paths of length ≥ 3). We tacitly identify T with its vertex set, which is assumed to be infinite. We write $u \sim v$ if $u, v \in T$ are neighbors. For any pair of vertices $u, v \in T$, there is a unique shortest path, called *geodesic segment*

$$\pi(u,v) = [u = v_0\,, v_1\,, \dots, v_k = v]$$

such that $v_{i-1} \sim v_i$ and all v_i are disctinct. If $u = v$ then this is the *empty* or *trivial* path. The number k is the *length* of the path (the graph distance between u and v). In T we choose and fix a *root vertex* o. We write $|v|$ for the length of $\pi(o,v)$. The choice of the root induces a partial order on T, where $u \leq v$ when $u \in \pi(o,v)$. Every $v \in T \setminus \{o\}$ has a unique *predecessor* v^-, which is the unique neighbour of v on $\pi(o,v)$. Thus, the set of all (unoriented) edges of T is

$$E(T) = \{[v^-, v] : v \in T\,,\ v \neq o\}\,.$$

For $u \in T$, the elements of the set

$$\{v \in T : v^- = u\}$$

are the *successors* of u, and its cardinality $\deg^+(u)$ is the *forward degree* of u.

In this and the next section, we assume that

$$2 \leq \deg^+(u) < \infty \quad \text{for every } \ u \in T\,. \tag{10.1}$$

A *(geodesic) ray* in T is a one-sided infinite path $\pi = [v_0\,, v_1\,, v_2\,, \dots]$ such that $v_{n-1} \sim v_n$ and all v_n are disctinct. Two rays are *equivalent* if their symmetric difference (as sets of vertices) is finite. An *end* of T is an equivalence class of rays. We shall typically use letters x, y, z to denote ends (and letters u, v, w for vertices). The set of all ends of T is denoted ∂T. This is the *boundary* at infinity of the tree. For any $u \in T$ and $x \in \partial T$, there is a unique ray $\pi(u,x)$ which is a representative of the end (equivalence class) x and starts at u. We write

$$\widehat{T} = T \cup \partial T.$$

For $u \in T$, the *branch of T rooted at u* is the subtree T_u that we identify with its set of vertices

$$T_u = \{v \in T : u \leq v\}\,,$$

so that $T_o = T$. We write ∂T_u for the set of all ends of T which have a representative path contained in T_u, and $\widehat{T}_u = T_u \cup \partial T_u$.

For $w, z \in \widehat{T}$, we define their *confluent* $w \wedge z = w \wedge_o z$ with respect to the root o by the relation

$$\pi(o, w \wedge z) = \pi(o,w) \cap \pi(o,z)\,.$$

It is the last common element on the geodesics $\pi(o,w)$ and $\pi(o,z)$, a vertex of T unless $w = z \in \partial T$.

One of the most common ways to define an ultra-metric on $\widehat{T}$ is

$$d_e(z,w) = \begin{cases} 0\,, & \text{if } z = w\,,\\ e^{-|z\wedge w|}\,, & \text{if } z \neq w\,. \end{cases} \tag{10.2}$$

Then $\widehat{T}$ is compact, and T is open and dense. We are mostly interested in the compact ultra-metric space ∂T. In the metric d_e of (10.2), each d_e-ball with centre $x \in \partial T$ is of the form ∂T_u for some $x \in \pi(o,x)$. Indeed

$$\partial T_u = B_{e^{-|u|}}(x) \quad \text{for every } o \in \pi(o,x)\,, \quad \text{and} \quad \Lambda_{d_e}(x) = \{e^{-|u|} : u \in \pi(o,u)\}\,.$$

Conversely, we now start with a compact ultra-metric space (X, d) that does not possess isolated points, and construct a tree T as follows: The vertex set of T is the collection

$$\mathcal{B} = \{B_r(x) : x \in X\ ,\ r > 0\}$$

of all closed balls in (X, d), already encountered in §7. Here, we may assume (if we wish) that $r \in \Lambda_d(x)$.

We now consider any ball $v = B \in \mathcal{B}$ as a vertex of a tree T. We choose our root vertex as $o = X$, which belongs to $\mathcal{B}$ by compactness. Neighborhood is given by the predecessor relation of balls, as given by Definition 2.1. That is, if $v = B$ then $u = B'$ is the predecessor vertex v^- of v in the tree T. By compactness, each x has only finitely many successors, and since there are no isolated points in X, every vertex has at least 2 successors, so that (10.1) holds.

This defines the tree structure. For any $x \in X$, the collection of all balls $B_r(x)$, $r \in \Lambda_d(x)$, ordered decreasingly, forms the set of vertices of a ray in T that starts at o. Via a straightforward exercise, the mapping that associates to x the end of T represented by that ray is a homeomorphism from X onto ∂T. Thus, we can identify X and ∂T as ultra-metric spaces.

In this identification, if originally a vertex u was interpreted as a ball $B_r(x)$, $r \in \Lambda_d(x)$, then the set ∂T_u of ends of the branch T_u just coincides with the ball $B_r(x)$. That is, we are identifying each vertex u of T with the set ∂T_u.

If we start with an arbitrary locally finite tree and take its space of ends as the ultra-metric space X, then the above construction does not recover vertices with forward degree 1, so that in general we do not get back the tree we started with. However, via the above construction, the correspondence between compact ultra-metric spaces without isolated points (perfect ultra-metric spaces) and locally finite rooted trees with forward degrees ≥ 2 is bijective.

It is well known that any ultra-metric space X which is both compact and perfect is homeomorphic to the ternary Cantor set $C \subset [0, 1]$. When X is not compact but still perfect we have a homeomorphism $X \simeq C \setminus \{p\}$, where $p \in C$ is any fixed point.

For the rest of this and the next section, we shall abandon the notation X for compact and perfect ultra-metric space.

We consider X as the boundary ∂T of a locally finite, rooted tree with forward degrees ≥ 2.

At the end, we shall comment on how one can handle the presence of vertices with forward degree 1, as well as the non-compact case.

There are many ways to equip ∂T with an ultra-metric that has the same topology and the same compact-open balls ∂T_x , $x \in T$, possibly with different radii than in the standard metric (10.2). The following is a kind of ultrametric analogue of a length element.

Definition 10.1 Let T be a locally finite, rooted tree T with $\deg^+(x) \geq 2$ for all x. An *ultra-metric element* is a function $\phi : T \to (0\,,\infty)$ with

$$\text{(i)}\quad \phi(v^-) > \phi(v) \quad \text{for every } v \in T \setminus \{o\}\,,$$
$$\text{(ii)}\quad \lim \phi(v_n) = 0 \quad \text{along every geodesic ray } \pi = [v_0\,, v_1\,, v_2\,, \dots]\,.$$

It induces the ultra-metric d_ϕ on ∂T given by

$$d_\phi(x, y) = \begin{cases} 0\,, & \text{if } x = y\,, \\ \phi(x \wedge y)\,, & \text{if } x \neq y\,. \end{cases}$$

The balls in this ultra-metric are again the sets

$$\partial T_u = B_{\phi(u)}(x)\,, \quad x \in \partial T_u\,.$$

Note that condition (ii) in the definition is needed for having that each end of T is non-isolated in the metric d_ϕ . The metric d_e of (10.2) is of course induced by $\phi(x) = e^{-|x|}$.

Lemma 10.2 *For a tree as in Definition 10.1, every ultra-metric on ∂T whose closed balls are the sets ∂T_u , $u \in T$, is induced by an ultra-metric element on T.*

Proof. Given an ultra-metric d as stated, we set $\phi(v) = \operatorname{diam}(\partial T_v)$, the diameter with respect to the metric d. Since $\deg^+(v^-) \ge 2$ for any $v \in T \setminus \{o\}$, the ball ∂T_{v^-} is the disjoint union of at least two balls ∂T_u with $u^- = v^-$. Therefore we must have $\operatorname{diam}(\partial T_v) < \operatorname{diam}(\partial T_{v^-})$, and property (i) holds. Since no end is isolated, ϕ satisfies (ii). It is now straightforward that $d_\phi = d$. ■

In view of this correspondence, in the sequel we shall replace the subscript d referring to the metric $d = d_\phi$ by the subscript ϕ referring to the ultra-metric element. We note that

$$\operatorname{diam}_\phi(\partial T) = \phi(o)\,, \quad \Lambda_\phi(x) = \{\phi(u) : u \in \pi(o,x)\} \quad \text{and} \quad \Lambda_\phi = \{\phi(v) : v \in T\}. \tag{10.3}$$

We also note here that for any $x \in \partial T$ and $v \in \pi(o,x)$, the balls with respect to d_ϕ are

$$B_r(x) = B_r^\phi(x) = \begin{cases} \partial T_v & \text{for } \phi(v) \le r < \phi(v^-)\,, \text{ if } v \ne o \\ \partial T & \text{for } r \ge \phi(o)\,, \text{ if } v = o\,. \end{cases} \tag{10.4}$$

Isotropic jump processes on the boundary of a tree. In view of the explanations given above, we can consider the isotropic jump processes of (1.2)–(1.4) on $X = \partial T$. Since this space is compact, we may assume that the reference measure μ is a probability measure on ∂T. Given μ, a distance distribution σ with properties (2.3), and an ultrametric element ϕ on T, we can now refer to the (d_ϕ, μ, σ)-process simply as the (ϕ, μ, σ)-process on ∂T. We can write the semigroup and its transition probabilities in detail as follows. For $x \in \partial T$ and $\pi(0,x) = [o = v_0\,, v_1\,, v_2\,, \dots]$, using (10.4),

$$P^t f(x) = \sum_{n=0}^{\infty} c_n^t \, P_{\phi(v_n)} f(x)\,,$$
$$\text{where} \quad c_0^t = 1 - \sigma^t\big(\phi(v_0)\big) \quad \text{and} \quad c_n^t = \sigma^t\big(\phi(v_{n-1})\big) - \sigma^t\big(\phi(v_n)\big) \;\; \text{for } n \ge 1\,.$$

Thus, for arbitrary $u \in T$ and $x \in \partial T$ as above

$$\mathbb{P}[X_t \in \partial T_u \mid X_0 = x] = \sum_{n=0}^{\infty} c_n^t \, \frac{\mu(\partial T_{v_n} \cap \partial T_u)}{\mu(\partial T_{v_n})}. \tag{10.5}$$

We know that we have some freedom in the choice of the measure σ: any two measures whose distribution functions coincide on the value set Λ_ϕ of ϕ give rise to the same process. Recall the Definition 1.3 of the standard (d, μ)-process, now to be re-named the standard (ϕ, μ)-process.

Nearest neighbour random walks on a tree. On a tree as a discrete structure, there are other, very well studied stochastic processes, namely *random walks.* Our aim is to analyze how they are related with isotropic jump processes on the boundary of the tree. A good part of the material outlined next is taken from the book of Woess [47]. An older, recommended reference is the seminal paper of Cartier [11].

A *nearest neighbour random walk* on the locally finite, infinite tree T is induced by its stochastic transition matrix $\mathcal{P} = \big(p(u,v)\big)_{u,v \in T}$ with the property that $p(u,v) > 0$ if and only if $u \sim v$. The resulting discrete-time Markov chain (random walk) is written $(Z_n)_{n \ge 0}$. Its n-step transition probabilities

$$p^{(n)}(u,v) = \mathbb{P}_u[Z_n = v], \quad u, v \in T,$$

are the elements of the n^{th} power of the matrix $\mathcal{P}$. The notation $\mathbb{P}_u$ refers to the probability measure on the trajectory space that governs the random walk starting at u. We assume that the random walk is *transient,* i.e., with probability 1 it visits any finite set only finitely often. Thus, $0 < G(u,v) < \infty$ for all $u, v \in T$, where

$$G(u,v) = \sum_{n=0}^{\infty} p^{(n)}(u,v)$$

is the *Green kernel* of the random walk. In addition, we shall also make crucial use of the quantities

$$F(u,v) = \mathbb{P}_u[Z_n = v \text{ for some } n \geq 0] \quad \text{and} \quad U(v,v) = \mathbb{P}_v[Z_n = v \text{ for some } n \geq 1]\,.$$

We shall need several identities relating them and start with a few of them, valid for all $u, v \in T$.

$$G(u,v) = F(u,v)G(v,v) \tag{10.6}$$

$$G(v,v) = \frac{1}{1 - U(v,v)} \tag{10.7}$$

$$U(v,v) = \sum_u p(v,u)F(u,v) \tag{10.8}$$

$$F(u,v) = F(u,w)F(w,v) \quad \text{whenever} \;\; w \in \pi(u,v) \tag{10.9}$$

The first three hold for arbitrary denumerable Markov chains, while (10.9) is specific for trees (resp., a bit more generally, when w is a so-called cut point between u and v). The identities show that those quantities are completely determined just by all the $F(u,v)$, where $u \sim v$. More identities, as to be found in [47, Chapter 9], will be displayed and used later on. By transience, the random walk Z_n must converge to a random end, a simple and well-known fact. See e.g. [11] or [47, Theorem 9.18].

Lemma 10.3 *There is a ∂T-valued random variable Z_∞ such that for every starting point $u \in T$,*

$$\mathbb{P}_u[Z_n \to Z_\infty \textit{ in the topology of } \widehat{T}] = 1.$$

In brief, the argument is as follows: by transience, random walk trajectories must accumulate at ∂T almost surely. If such a trajectory had two distinct accumulation points, say x and y, then by the nearest neighbour property, the trajectory would visit the vertex $x \wedge y$ infinitely often, which can occur only with probability 0.

We can consider the family of *limit distributions* ν_u , $u \in T$, where for any Borel set $B \subset \partial T$,

$$\nu_u(B) = \mathbb{P}_u[Z_\infty \in B]\,.$$

The sets ∂T_u , $u \in T$ (plus the empty set), form a semi-algebra that generates the Borel σ-algebra of ∂T. Thus, each ν_u is determined by the values of those sets. There is an explicit formula, compare with [11] or [47, Proposition 9.23]. For $v \neq o$,

$$\nu_u(\partial T_v) = \begin{cases} F(u,v)\dfrac{1 - F(v,v^-)}{1 - F(v^-,v)F(v,v^-)}\,, & \text{if } u \in \{v\} \cup (T \setminus T_v)\,, \\[2ex] 1 - F(u,v)\dfrac{F(v,v^-) - F(v^-,v)F(v,v^-)}{1 - F(v^-,\nu)F(v,v^-)}\,, & \text{if } u \in T_v\,. \end{cases} \tag{10.10}$$

A *harmonic function* is a function $h : T \to \mathbb{R}$ with $\mathcal{P}h = h$, where

$$Ph(u) = \sum_v p(u,v)h(v)\,.$$

For any Borel set $B \subset \partial T$, the function $u \mapsto \nu_u(B)$ is a bounded harmonic function. One deduces that all ν_u are comparable: $p^{(k)}(u,v)\,\nu_u \le \nu_v$, where k is the length of $\pi(u,v)$. Thus, for any function $\varphi \in L^1(\partial T, \nu_o)$, the function h_φ defined by

$$h_\varphi(u) = \int_{\partial T} \varphi \, d\nu_u$$

is finite and harmonic on T. It is often called the *Poisson transform* of φ.

We next define a measure m on T via its atoms: $\mathsf{m}(o) = 1$, and for $v \in T \setminus \{o\}$ with $\pi(o,v) = [o = v_0\,, v_1\,, \dots, v_k = v]$,

$$\mathsf{m}(v) = \frac{p(v_0,v_1)p(v_1,v_2)\cdots p(v_{k-1},v_k)}{p(v_1,v_0)p(v_2,v_1)\cdots p(v_k,v_{k-1})}\,. \tag{10.11}$$

Then for all $u, v \in T$,

$$\mathsf{m}(u)p(u,v) = \mathsf{m}(v)p(v,u)\,, \quad \text{and consequently} \quad \mathsf{m}(u)G(u,v) = \mathsf{m}(v)G(v,u)\,; \tag{10.12}$$

the random walk is *reversible*. This would allow us to use the *electrical network* interpretation of $(T, \mathcal{P}, \mathsf{m})$, for which there are various references: see e.g. Yamasaki [48], Soardi [39], or – with notation as used here – [47, Chapter 4]. We do not go into its details here; each edge $e = [v^-, v] \in E(T)$ is thought of as an electric conductor with *conductance*

$$a(v^-, v) = \mathsf{m}(v)p(v, v^-).$$

We get the Dirichlet form $\mathcal{E}_T = \mathcal{E}_{T,\mathcal{P}}$ for functions $f, g : T \to \mathbb{R}$, defined by

$$\mathcal{E}_T(f,g) = \sum_{[v^-,v] \in E(T)} \big(f(v) - f(v^-)\big)\big(g(v) - g(v^-)\big)\, a(v^-, v)\,. \tag{10.13}$$

It is well defined for f, g in the space

$$\mathcal{D}(T) = \mathcal{D}(T, \mathcal{P}) = \{f : T \to \mathbb{R} \mid \mathcal{E}_T(f,f) < \infty\}. \tag{10.14}$$

Harmonic functions of finite energy and their boundary values. We are interested in the subspace

$$\mathcal{HD}(T) = \mathcal{HD}(T, \mathcal{P}) = \{h \in \mathcal{D}(T, \mathcal{P}) : \mathcal{P}h = h\}$$

of harmonic functions with *finite power.* The terminology comes from the interpretation of such a function as the potential of an electric flow (or current), and then $\mathcal{E}_T(h,h)$ is the power of that flow.[2]

Every function in $\mathcal{HD}(T, \mathcal{P})$ is the Poisson transform of some function $\varphi \in L^2(\partial T, \nu_o)$. This is valid not only for trees, but for general finite range reversible Markov chains, and follows from the following facts.

1. Every function in $\mathcal{HD}$ is the difference of two non-negative functions in $\mathcal{HD}$.
2. Every non-negative function in $\mathcal{HD}$ can be approximated, monotonically from below, by a sequence of non-negative bounded functions in $\mathcal{HD}$.
3. Every bounded harmonic function (not necessarily with finite power) is the Poisson transform of a bounded function on the boundary.

---

[2] In the mathematical literature, mostly the expression "energy" is used for $\mathcal{E}_T(h,h)$, but it seems that "power" is the more appropriate terminology from Physics.

In the general setting, the latter is the (active part of) the Martin boundary, with ν_u being the limit distribution of the Markov chain, starting from u, on that boundary. (1) and (2) are contained in [48] and [39], while (3) is part of general Martin boundary theory, see e.g. [47, Theorem 7.61].

Thus, we can introduce a form $\mathcal{E}_{\mathcal{HD}}$ on ∂T by setting

$$\begin{aligned} \mathcal{D}(\partial T, \mathcal{P}) &= \{\varphi \in L^1(\partial T, \nu_o) : \mathcal{E}_T(h_\varphi, h_\varphi) < \infty\}, \\ \mathcal{E}_{\mathcal{HD}}(\varphi, \psi) &= \mathcal{E}_T(h_\varphi, h_\psi) \quad \text{for } \varphi, \psi \in \mathcal{D}(\partial T, \mathcal{P}). \end{aligned} \tag{10.15}$$

Jump processes on the boundary of a tree. Kigami [24] elaborates an expression for the form $\mathcal{E}_{\mathcal{HD}}(\varphi, \psi)$ of (10.15) by considerable effort, shows its regularity properties and then studies the jump process on ∂T induced by this Dirichlet form. We call this the *boundary process* associated with the random walk.

Now, there is a rather simple expression for $\mathcal{E}_{\mathcal{HD}}$. We define the *Naïm kernel* on $\partial T \times \partial T$ by

$$\Theta_o(x, y) = \begin{cases} \dfrac{\mathsf{m}(o)}{G(o,o)F(o, x \wedge y)F(x \wedge y, o)}, & \text{if } x \neq y, \\ +\infty, & \text{if } x = y. \end{cases} \tag{10.16}$$

In our case, $\mathsf{m}(o) = 1$, but we might want to change the base point, or normalize the measure m in a different way.

Theorem 10.4 *For any transient nearest neighbour random walk on the tree T with root o, and all functions φ, ψ in $\mathcal{D}(\partial T, \mathcal{P})$,*

$$\mathcal{E}_{\mathcal{HD}}(\varphi, \psi) = \frac{1}{2} \int_{\partial T} \int_{\partial T} \big(\varphi(x) - \varphi(y)\big)\big(\psi(x) - \psi(y)\big) \Theta_o(x, y)\, d\nu_o(x)\, d\nu_o(y).$$

There is a general definition of the Naïm kernel [30] that involves the Martin boundary, which in the present case is ∂T. A proof of Theorem 10.4 is given in [16] in a setting of abstract potential theory on Green spaces, which are locally Euclidean. The definition of [30] refers to the same type of setting. Now, infinite networks, even when seen as metric graphs, are not locally Euclidean. In this sense, so far the definition of the kernel and a proof of Theorem (10.4) for transient, reversible random walks have not been well accessible in the literature. In a forthcoming paper, Georgakopoulos and Kaimanovich [19] will provide those "missing links". We shall give a direct and simple proof of Theorem 10.4 for the specific case of trees. We start with the following observation.

Lemma 10.5 *The measure $\Theta_o(x, y)\, d\nu_o(x)\, d\nu_o(y)$ on $\partial T \times \partial T$ is invariant with respect to changing the base point (root) o.*

Proof. We want to replace the base point o with some other $u \in T$. We may assume that $u \sim o$. Indeed, then we may step by step replace the current base point by one of its neighbors to obtain the result for arbitrary u.

Recall that the confluent that appears in the definition (10.16)) of Θ_o depends on the root o, while for Θ_x it becomes the one with respect to x as the new root. It is a well-known fact that

$$\frac{d\nu_u}{d\nu_o}(x) = K(u, x) = \frac{G(u, u \wedge_o x)}{G(o, u \wedge_o x)},$$

the *Martin kernel.* Thus, we have to show that for all $x, y \in \partial T$ $(x \neq y)$

$$\frac{\mathsf{m}(o)}{G(o,o)F(o, x \wedge_o y)F(x \wedge_o y, o)} = \frac{\mathsf{m}(u)K(u,x)K(u,y)}{G(u,u)F(u, x \wedge_u y)F(x \wedge_u y, u)}.$$

Case 1. $x, y \in \partial T_u$. Then $x \wedge_o y = x \wedge_u y =: v \in T_u$, and $u \wedge_o x = u \wedge_o y = u$. Thus, using (10.6), (10.9) and the fact that by (10.12)

$$\mathsf{m}(u)/G(o,u) = \mathsf{m}(o)/G(u,o),$$

we obtain

$$\begin{aligned}
\frac{\mathsf{m}(u)K(u,x)K(u,y)}{G(u,u)F(u,x\wedge_u y)F(x\wedge_u y,u)} &= \frac{\mathsf{m}(u)}{G(u,u)F(u,v)F(v,u)}\left(\frac{G(u,u)}{G(o,u)}\right)^2 \\
&= \frac{\mathsf{m}(o)G(u,u)}{F(u,v)F(v,u)G(o,u)G(u,o)} \\
&= \frac{\mathsf{m}(o)}{F(u,v)F(v,u)F(o,u)F(u,o)G(o,o)} \\
&= \frac{\mathsf{m}(o)}{F(o,v)F(v,o)G(o,o)},
\end{aligned}$$

as required. There are 3 more cases.

Case 2. $x, y \in \partial T \setminus \partial T_u$. Then

$$x \wedge_o y = x \wedge_u y =: w \in T \setminus T_u, \quad \text{and} \quad u \wedge_o x = u \wedge_o y = o.$$

Case 3. $x \in \partial T_u$, $y \in \partial T \setminus \partial T_u$. Then

$$x \wedge_o y = o, \quad x \wedge_u y = u, \quad u \wedge_o x = u \quad \text{and} \quad u \wedge_o y = o.$$

Case 4. $x \in \partial T \setminus \partial T_u$, $y \in \partial T_u$. This is like Case 3, exchanging the roles of x and y.

In all cases 2–4, the computation is done very similarly to Case 1, a straightforward exercise. ■

For proving Theorem 10.4, we need a few more facts related with the network setting; compare e.g. with [47, §4.D].

The space $\mathcal{D}(T)$ of (10.14) is a Hilbert space when equipped with the inner product

$$(f,g) = \mathcal{E}_T(f,g) + f(o)g(o).$$

The subspace $\mathcal{D}_0(T)$ is defined as the closure of the space of finitely supported functions in $\mathcal{D}(T)$. It is a proper subspace if and only if the random walk is transient, and then the function $G_v(u) = G(u,v)$ is in $\mathcal{D}_0(T)$ for any $v \in T$ [48], [39]. We need the formula

$$\mathcal{E}_T(f, G_v) = \mathsf{m}(v)f(v) \quad \text{for every } f \in \mathcal{D}_0(T). \tag{10.17}$$

Given a branch T_w of T ($w \in T \setminus \{o\}$), we can consider it as a subnetwork equipped with the same conductances $a(u,v)$ for $[u,v] \in E(T_w)$. The associated measure on T_w is

$$\mathsf{m}_{T_w}(u) = \sum_{v \in T_z : v \sim u} a(u,v) = \begin{cases} \mathsf{m}(u) & \text{if } u \in T_w \setminus \{w\}, \\ \mathsf{m}(w) - a(w,w^-) & \text{if } u = w. \end{cases}$$

The resulting random walk on T_w has transition probabilities

$$p_{T_w}(u,v) = \frac{a(v,w)}{\mathsf{m}_{T_w}(u)} = \begin{cases} p(u,v) & \text{if } u \in T_w \setminus \{w\},\ v \sim u, \\[2ex] \dfrac{p(w,v)}{1 - p(w,w^-)} & \text{if } u = w,\ v \sim u. \end{cases}$$

We have $F_{T_w}(u,u^-) = F(u,u^-)$ and thus also $F_{T_w}(u,w) = F(u,w)$ for every $u \in T_w \setminus \{w\}$, because before its first visit to w, the random walk on T_w obeys the same transition probabilities as the original random walk on T. It is then easy to see [47, p. 241] that the random walk on T_w is transient if and only if for the original random walk, $F(w,w^-) < 1$, which in turn holds if and only if $\nu_o(\partial T_w) > 0$. (In other parts of this and the preceding two sections, this is always assumed, but for the proof of Theorem 10.4, we just assume the random walk on the whole of T to be transient.) Conversely, if $F(w,w^-) = 1$ then $F(u,w) = 1$ for all $u \in T_w$.

Below, we shall need the following formula for the limit distributions.

Lemma 10.6 *For* $u \in T \setminus \{o\}$,

$$\nu_v(\partial T_u) = 1 - p(u,u^-)\big(G(u,u) - G(u^-,u)\big)\,.$$

Proof. By (11.7),

$$G(u,u)p(u,u^-) = \frac{F(u,u^-)}{1 - F(u,u^-)F(u^-,u)}$$

Thus,

$$p(u,u^-)\big(G(u,u) - G(u^-,u)\big) = \big(1 - F(u^-,u)\big)G(u,u)p(u,u^-) = 1 - \nu_u(\partial T_u)$$

after a short computation using (10.10) ∎

Proof of Theorem 10.4. We first prove the Doob- Naïm formula (shortly, D-N-formula) for the case when $\varphi = \mathbf{1}_{\partial T_v}$ and $\psi = \mathbf{1}_{\partial T_w}$ for two proper branches T_v and T_w of T. They are either disjoint, or one of them contains the other.

Case 1. $T_w \subset T_v$. (The case $T_v \subset T_w$ is analogous by symmetry.)

This means that $w \in T_v$. For $x, y \in \partial T$ we have

$$\big(\varphi(x) - \varphi(y)\big)\big(\psi(x) - \psi(y)\big) = 1$$

if $x \in \partial T_w$ and $y \in \partial T \setminus \partial T_v$ or conversely, and $= 0$ otherwise. By Lemma 10.5, we may choose v as the base point. Thus, the right hand side of the identity is

$$\int_{\partial T \setminus \partial T_v} \int_{\partial T_w} \Theta_v(x,y)\, d\nu_v(x)\, d\nu_v(y) = \frac{\mathsf{m}(v)}{G(v,v)}\, \nu_v(\partial T \setminus \partial T_v)\nu_v(\partial T_w)\,,$$

since $x \wedge_v y = v$ and $F(v,v) = 1$.

Let us now turn to the left hand side of the D-N-formula. The Poisson transforms of φ and ψ are

$$h_\varphi(u) = \nu_u(\partial T_v) \quad \text{and} \quad h_\psi(u) = \nu_u(\partial T_w).$$

By (10.10),

$$\begin{aligned} h_\varphi(u) &= F(u,v)\nu_v(\partial T_v)\,, \quad u \in \{v\} \cup (T \setminus T_v) \\ 1 - h_\varphi(u) &= F(u,v)(\partial T \setminus \partial T_v)\,, \quad u \in T_v\,. \end{aligned}$$

We set $F_v(u) = F(u,v)$ and write

$$h_\varphi(u) - h_\varphi(u^-) = \big(1 - h_\varphi(u^-)\big) - \big(1 - h_\varphi(u)\big)$$

whenever this is convenient, and analogously for h_ψ . Then we get

$$\begin{aligned}
&\mathcal{E}_T(h_\varphi\,,h_\psi)\\
&= \sum_{[u,u^-]\in E(T)\setminus E(T_v)} a(u,u^-)\big(F(u,v)-F(u^-,v)\big)\nu_v(\partial T_v)\,\big(F(u,v)-F(u^-,v)\big)\nu_v(\partial T_w)\\
&\quad - \sum_{[u,u^-]\in E(T_v)\setminus E(T_w)} a(u,u^-)\big(F(u,v)-F(u^-,v)\big)\nu_v(\partial T\setminus\partial T_v)\,\big(F(u,v)-F(u^-,v)\big)\nu_v(\partial T_w)\\
&\quad + \sum_{[u,u^-]\in E(T_w)} a(u,u^-)\big(F(u,v)-F(u^-,v)\big)\nu_v(\partial T\setminus\partial T_v)\,\big(F(u,v)-F(u^-,v)\big)\nu_v(\partial T\setminus\partial T_w)\\
&= \mathcal{E}_T(F_v\,,F_w)\nu_v(\partial T_v)\nu_v(\partial T_w) - \mathcal{E}_{T_v}(F_v\,,F_w)\nu_v(\partial T_w) + \mathcal{E}_{T_w}(F_v\,,F_w)\nu_v(\partial T\setminus\partial T_v)\,,
\end{aligned}$$

where of course $\mathcal{E}_{T_v}$ is the Dirichlet form of the random walk on the branch T_v , as discussed above, and analogously for $\mathcal{E}_{T_w}$. Now $F_v = G_v/G(v,v)$ by (10.6), whence (10.17) yields

$$\mathcal{E}_T(F_v,F_w) = \frac{\mathcal{E}_T(G_v,F_w)}{G(v,v)} = \frac{\mathsf{m}(v)F(v,w)}{G(v,v)}\,. \tag{10.18}$$

Recall that for the random walk on T_v , we have $F_{T_v}(u,v) = F(u,v)$ for every $u\in T_v$. Also,

$$\mathsf{m}_{T_v}(v) = \mathsf{m}(v) - a(v,v^-) = \mathsf{m}(v)\big(1-p(v,v^-)\big).$$

We apply (10.18) to that random walk and get

$$\mathcal{E}_{T_v}(F_v,F_w) = \frac{\mathsf{m}(v)\big(1-p(v,v^-)\big)F(v,w)}{G_{T_v}(v,v)}\,.$$

We now apply (10.7) and (10.8), recalling in addition that

$$p_{T_v}(v,u) = \frac{p(v,u)}{1-p(v,v^-)}$$

for $u\in T_v$, and obtain

$$\begin{aligned}
\frac{1-p(v,v^-)}{G_{T_v}(v,v)} &= 1-p(v,v^-) - \big(1-p(v,v^-)\big)U_{T_v}(v,v)\\
&= 1-p(v,v^-) - \sum_{u:u^-=v} p(v,u)F(u,v)\\
&= 1-p(v,v^-) - \big(U(v,v)-p(v,v^-)F(v^-,v)\big)\\
&= \frac{1}{G(v,v)} - p(v,v^-)\big(1-F(v^-,v)\big) = \frac{\nu_v(\partial T_v)}{G(v,v)},
\end{aligned}$$

where in the last step we have used Lemma 10.6. We have obtained

$$\mathcal{E}_{T_v}(F_v,F_w) = \frac{\mathsf{m}(v)F(v,w)}{G(v,v)}\,\nu_v(\partial T_v).$$

In the same way, exchanging roles between T_w and T_v and using reversibility (10.12),

$$\mathcal{E}_{T_w}(F_v,F_w) = \frac{\mathsf{m}(w)F(w,v)}{G(w,w)}\,\nu_w(\partial T_w) = \frac{\mathsf{m}(v)F(v,w)}{G(v,v)}\,\nu_w(\partial T_w) = \frac{\mathsf{m}(v)}{G(v,v)}\,\nu_v(\partial T_w)$$

Putting things together, we get that

$$\mathcal{E}_T(h_\varphi\,,h_\psi) = \mathcal{E}_{T_w}(F_v\,,F_w)\nu_v(\partial T\setminus\partial T_v) = \frac{\mathsf{m}(v)}{G(v,v)}\,\nu_v(\partial T_w)\nu_v(\partial T\setminus\partial T_v),$$

as proposed.

Case 2. $T_w \cap T_v = \emptyset$.

In view of Lemma 10.5, both sides of the D-N-formula are independent of the root o. Thus we may declare our root to be one of the neighbors of v that is not on $\pi(v,w)$. Also, let $\bar{v}$ be the neighbour of v on $\pi(w,v)$. Then, with our chosen new root, the complement of the "old" T_v is $T_{\bar{v}}$, which contains T_w (The latter remains the same with respect to the new root).

Thus, we can apply the result of case 1 to $T_{\bar{v}}$ and T_w. This means that we have to replace the functions φ and h_φ with $1-\varphi$ and $1-h_\varphi$, respectively, which just means that we change the sign on both sides of the identity. We are re-conducted to Case 1 without further computations.

We deduce from what we have done so far, and from linearity of the Poisson transform as well of bilinearity of the forms on both sides of the D-N-formula , that it holds for linear combinations of indicator functions of sets ∂T_v. Those indicator functions are dense in the space $C(\partial T)$ with respect to the max-norm. Thus, the D-N-formula holds for all continuous functions on ∂T. The extension to all of $\mathcal{D}(\partial T, \mathcal{P})$ is by standard approximation. ■

11 Duality of random walks on trees and isotropic processes on their boundaries

When looking at our isotropic processes and at the *boundary process* of Kigami [24], it is natural to ask the following two questions.

Question I. Given a transient random walk on T associated with the Dirichlet form $\mathcal{E}_T$ of (10.13), does the boundary process on ∂T induced by the form $\mathcal{E}_{\mathcal{HD}}$ of (10.15) arise as one of the isotropic processes (1.4) on ∂T with transition probabilities (10.5), with respect to the measure $\mu = \nu_o$ on ∂T, some ultra-metric element ϕ on T and a suitable distance distribution σ on $[0\,,\,\infty)$?

Question II. Conversely, given data μ, ϕ and σ, is there a random walk on T with limit distribution $\nu_o = \mu$ such that the isotropic process induced by μ, ϕ and σ is the boundary process with Dirichlet form $\mathcal{E}_{\mathcal{HD}}$?

Before answering both questions, we need to specify the assumptions more precisely. When starting with (ϕ, μ, σ), we assume as in §1 that μ is supported by the whole of ∂T.

Thus, on the side of the random walk, we also want that $\operatorname{supp}(\nu_o) = \partial T$. This is equivalent with the requirement that $\nu_o(\partial T_v) > 0$ for every $v \in T$. By (10.10) this is in turn equivalent with

$$F(v, v^-) < 1 \quad \text{for every } v \in T \setminus \{o\}. \tag{11.1}$$

Indeed, we shall see that we need a bit more, namely that

$$\lim_{v \to \infty} G(v, o) = 0\,, \tag{11.2}$$

that is, for every $\varepsilon > 0$ there is a finite set $A \subset T$ such that $G(v, o) < \varepsilon$ for all $v \in T \setminus A$. This condition is necessary and sufficient for solvability of the *Dirichlet problem:* for any $\varphi \in C(\partial X)$, its Poisson transform h_φ provides the unique continuous extension of φ to $\widehat{T}$ which is harmonic in T. See e.g. [47, Corollary 9.44].

We shall restrict attention to random walks with properties (11.1) and (11.2) on a rooted tree with forward degrees ≥ 2.

Answer to Question I. We start with a random walk that fulfills the above requirements. We know from §1 that each (μ, ϕ, σ)-process arises as the standard process of Definition 1.3 with respect to the intrinsic metric, see the lines before (1.6): given ϕ and σ, the intrinsic metric is induced by the ultrametric element

$$\phi_*(u) = -1\big/ \log \sigma\big(\phi(u)\big)\,. \tag{11.3}$$

Thus, we can eliminate σ from our considerations by just looking for an ultra-metric element ϕ such that the $\partial-$ process is the standard process on ∂T associated with $(\nu_0\,, \phi)$.

Since the processes are determined by the Dirichlet forms, we infer from Theorems 7.2 and 10.4 that we are looking for ϕ such that $J(x,y) = \Theta(x,y)$ for all $x, y \in \partial T$ with $x \neq y$, where $J(x,y)$ is given by (7.1). Rewriting $J(x,y)$ in terms of ϕ, ν_o and the tree structure, this becomes

$$\frac{1}{\phi(o)} + \int_{1/\phi(o)}^{1/\phi(x\wedge y)} \frac{dt}{\nu_o\big(B^{\phi}_{1/t}(x)\big)} = \frac{1}{G(o,o)F(o,x\wedge y)F(x\wedge y,o)}. \tag{11.4}$$

First of all, since $\deg^+(o) \geq 2$, there are $x, y \in \partial T$ such that $x \wedge y = o$. We insert these two boundary points in (11.4)). Since $F(o,o) = 1$, we see that we must have

$$\phi(o) = G(o,o)\,.$$

Now take $v \in T \setminus \{o\}$. Since forward degrees are ≥ 2, there are $x, y, y' \in \partial T$ such that $x \wedge y = v$ and $x \wedge y' = v^-$. We write (11.4) first for (x, y') and then for (x,y) and then take the difference, leading to the equation

$$\int_{1/\phi(v^-)}^{1/\phi(v)} \frac{dt}{\nu_o\big(B^{\phi}_{1/t}(x)\big)} = \frac{1}{G(o,o)F(o,v)F(v,o)} - \frac{1}{G(o,o)F(o,v^-)F(v^-,o)}. \tag{11.5}$$

By (10.4), within the range of the last integral we must have $B^{\phi}_{1/t}(v) = \partial T_v$, whence that integral reduces to

$$\left(\frac{1}{\phi(v)} - \frac{1}{\phi(v^-)}\right)\frac{1}{\nu_o(\partial T_v)}$$

We multiply equation (11.5) by $\nu_o(\partial T_v)$ and simplify the resulting right hand side

$$\left(\frac{1}{G(o,o)F(o,v)F(v,o)} - \frac{1}{G(o,o)F(o,v^-)F(v^-,o)}\right)\nu_o(\partial T_v)$$

by use of the identities (10.6) – (10.9)) and the first of the two formulas of (10.10) (for ν_o). We obtain that the ultra-metric element that we are looking for should satisfy

$$\frac{1}{\phi(v)} - \frac{1}{\phi(v^-)} = \frac{1}{G(v,o)} - \frac{1}{G(v^-,o)} \quad \text{for every } v \in T \setminus \{o\}\,. \tag{11.6}$$

This determines $1/\phi(v)$ recursively, and we get

$$\phi(v) = G(v,o)\,.$$

Since by (10.6) and (10.9)

$$G(v,o) = F(v,v^-)G(v^- o),$$

the assumptions (11.1) and (11.2) yield that ϕ is an ultra-metric element. Tracing back the last computations, we find that with this choice of ϕ, we have indeed that $J(x,y) = \Theta(x,y)$ for all $x, y \in \partial T$ with $x \neq y$. We have proved the following.

Theorem 11.1 *Let T be a locally finite, rooted tree with forward degrees ≥ 2. Consider a transient nearest neighbour random walk on T that satisfies (10.6) and (10.9). Then the boundary process on ∂T induced by the Dirichlet form (10.15) coincides with the standard process associated with ultra-metric element $\phi = G(\cdot, o)$ and the limit distribution ν_o of the random walk.*

Let $\mathcal{L}$ be the Laplacian associated with the boundary process of Theorem 11.1. $\mathcal{L}$ acts on locally constant functions f by

$$\mathcal{L}f(x) = \int\limits_{\partial T} (f(x) - f(y))\Theta_o(x,y)d\nu_o(y).$$

In view of the identification of balls in ∂T with vertices of T, the functions of (7.6) now become

$$f_v = \frac{\mathbf{1}_{\partial T_v}}{\nu_o(\partial T_v)} - \frac{\mathbf{1}_{\partial T_{v^-}}}{\nu_o(\partial T_{v^-})}, \ v \in T\backslash\{o\}.$$

In addition, we set $f_o = \mathbf{1}$ and note that it is an eigenfunction of $\mathcal{L}$ with eigenvalue 0. Applying Theorem 7.3 we obtain

Corollary 11.2 *For $v \in T\backslash\{o\}$, we have $\mathcal{L}f_v = G(v^-, o)^{-1} f_v$. In particular,*

$$\mathrm{Spec}(\mathcal{L}) = \{G(v,o)^{-1} : v \in T\} \cup \{0\}.$$

Remark 11.3 For any two vertices v and w in $T\backslash\{o\}$ such that $v^- = w^- = u$ the functions f_v and f_w are eigenfunctions of $\mathcal{L}$ corresponding to the eigenvalue $\lambda = 1/G(u,o)$. Hence the eigenspace $\mathcal{H}(u)$ corresponding to the vertex u is spanned by functions $\{f_v : v^- = u\}$. Since the rank of the system $\{f_v : v^- = u\}$ is $\deg^+(u) - 1$, where $\deg^+(u) \geq 2$ is the forward degree of the vertex u, we obtain

$$\dim \mathcal{H}(u) = \deg^+(u) - 1.$$

Remark 11.4 Given the random walk on T and the associated boundary process on ∂T, we might want to realize it as the (ν_o, ϕ, σ)-process for an ultra-metric element ϕ different from $G(\cdot, o)$. This means that we have to look for a suitable distance distribution σ on $[0\,,\infty)$, different from the inverse exponential distribution (1.5). In view of (11.3), we are looking for σ such that for our given generic ϕ,

$$\sigma\big(\phi(v)\big) = e^{-1/G(v,o)}.$$

For this it is necessary that $\phi(u) = \phi(v)$ whenever $G(u,o) = G(v,o)$: we need ϕ to be constant on equipotential sets. In that case, the distribution function $\sigma(r)$ is determined by the above equation for r in the value set Λ_ϕ of the ultra-metric d_ϕ. We can "interpolate" that function in an arbitrary way (monotone increasing, left continuous) and get a feasible measure σ.

Answer to Question II. Answering question (II) means that we start with ϕ and μ and then look for a random walk with limit distribution $\nu_o = \mu$ such that the standard (ϕ, μ)-process is the boundary process associated with the random walk. We know from Theorem 11.1 that in this case, we should have $\phi(v) = G(v,o)$, whence in particular, $\phi(o) > 1$. Thus we cannot expect that every ϕ is suitable. The most natural choice is to replace ϕ by $C \cdot \phi$ for some constant $C > 0$. For the standard processes associated with ϕ and $C \cdot \phi$, respectively, this just gives rise of a linear time change: if the old process is $\{X_t\}_{t>0}$, then the new one is $\{X_{t/C}\}_{t>0}$.

Theorem 11.5 *Let T be a locally finite, rooted tree with forward degrees ≥ 2. Consider an ultra-metric element ϕ on T and a fully supported probability measure μ on ∂T. Then there are a unique constant $C > 0$ and a unique transient nearest neighbour random walk on T that satisfies (10.6) and (10.9) with the following properties:*

1. $\mu = \nu_o$ is the limit distribution of the random walk.
2. The associated boundary process coincides with the standard process on ∂T induced by the ultra-metric element $C \cdot \phi$ and the given measure μ.

For the proof, we shall need three more formulas. The first two are taken from [47, Lemma 9.35], while the third is immediate from (10.10) and (10.9)

$$G(u,u)\,p(u,v) = \frac{F(u,v)}{1 - F(u,v)F(v,u)} \quad \text{if } u \sim v\,, \quad \text{and} \tag{11.7}$$

$$G(u,u) = 1 + \sum_{v:v\sim u} \frac{F(u,v)F(v,u)}{1 - F(u,v)F(v,u)} \tag{11.8}$$

$$F(v^-,v) = \frac{\nu_o(\partial T_v)/F(o,v^-)}{1 - F(v,v^-) + F(v,v^-)\ \nu_o(\partial T_v)/F(o,v^-)}. \tag{11.9}$$

Proof of Theorem 11.5. We proceed as follows: we start with ϕ and μ and replace ϕ by a new ultra-metric element $C \cdot \phi$, with C to be determined, and μ being the candidate for the limit distribution of the random walk that we are looking for.

Using the various formulas at our disposal, we first construct in the only possible way the quantities $F(u,v)$, $u,v,\in T$, in particular when $u \sim v$. In turn, they lead to the Green kernel $G(u,v)$. So far, these will be only "would-be" quantities whose feasibility will have to be verified. Until that verification, we shall denote them $\widetilde{F}(u,v)$ and $\widetilde{G}(u,v)$. Via (11.7), they will lead to definitions of transition probabilities $p(u,v)$. Stochasticity of the resulting transition matrix $\mathcal{P}$ will also have to be verified.

Only then, we will use a potential theoretic argument to show that $\widetilde{G}(u,v)$ really is the Green kernel associated with $\mathcal{P}$, so that the question mark that is implicit in the "$\widetilde{\ }$" symbol can be removed.

First of all, in view of Theorem 11.1, we must have

$$C \cdot \phi(v) = \widetilde{G}(v,o)\,,$$

whence by (10.6) and (10.9)

$$\widetilde{F}(v,v^-) = \phi(v)/\phi(v^-) \quad \text{for } v \in T \setminus \{o\}\,, \tag{11.10}$$

and more generally

$$\widetilde{F}(v,u) = \phi(v)/\phi(u) \quad \text{when } u \leq v\,.$$

We note immediately that $0 < \widetilde{F}(v,u) < 1$ when $u < v$, and that $\widetilde{F}(u,u) = 1$.

Next, we use (11.9) to construct recursively $\widetilde{F}(v^-,v)$ and $\widetilde{F}(o,v)$. We start with $\widetilde{F}(o,o) = 1$. If $v \neq o$ and $\widetilde{F}(o,v^-)$ is already given, with

$$\mu(\partial T_{v^-}) \leq \widetilde{F}(o,v^-) \leq 1$$

(the lower bound is required by (10.10)), then we have to set

$$\widetilde{F}(v^-,v) = \frac{\mu(\partial T_v)/\widetilde{F}(o,v^-)}{1 - \widetilde{F}(v,v^-) + \widetilde{F}(v,v^-)\ \mu(\partial T_v)/\widetilde{F}(o,v^-)} \tag{11.11}$$

and

$$\widetilde{F}(o,v) = \widetilde{F}(o,v^-)\widetilde{F}(v^-,v)\,.$$

Since

$$\widetilde{F}(o,v^-) \ge \mu(\partial T_{v^-}) \ge \mu(\partial T_v),$$

we see that

$$0 < \widetilde{F}(v^-,v) \le 1.$$

We set – as imposed by (10.9) –

$$\widetilde{F}(o,v) = \widetilde{F}(o,v^-)\widetilde{F}(v^-,v).$$

Formula (11.11) (re-)transforms into

$$\mu(\partial T_v) = \widetilde{F}(o,v^-)\widetilde{F}(v^-,v)\frac{1-\widetilde{F}(v,v^-)}{1-\widetilde{F}(v,v^-)\widetilde{F}(v^-,v)} \le \widetilde{F}(o,v) \le 1\,, \tag{11.12}$$

as needed for our recursive construction. At this point, we have all $\widetilde{F}(u,v)$, initially for $u \sim v$, and consequently for all u,v by taking products along geodesic paths.

We now can compute the constant C: (11.8), combined with (11.10) and (11.12) for $u \sim o$ forces

$$\begin{aligned}
C\phi(o) &= \widetilde{G}(o,o) = 1 + \sum_{u:u\sim o} \frac{\widetilde{F}(o,u)\widetilde{F}(u,o)}{1-\widetilde{F}(o,u)\widetilde{F}(u,o)} \\
&= 1 + \sum_{u:u\sim o} \frac{\widetilde{F}(u,o)}{1-\widetilde{F}(u,o)}\mu(\partial T_u) \\
&= 1 + \sum_{u:u\sim o} \frac{\phi(u)/\phi(o)}{1-\phi(u)/\phi(o)}\mu(\partial T_u)
\end{aligned}$$

Therefore

$$C = \frac{1}{\phi(o)} + \sum_{u:u\sim o} \frac{\phi(u)/\phi(o)}{\phi(o)-\phi(u)}\mu(\partial T_u)\,. \tag{11.13}$$

We now construct $\widetilde{G}(u,u)$ via (11.8):

$$\widetilde{G}(u,u) = 1 + \sum_{v:v\sim u} \frac{\widetilde{F}(u,v)\widetilde{F}(v,u)}{1-\widetilde{F}(u,v)\widetilde{F}(v,u)}. \tag{11.14}$$

For $u = o$, we know that this is compatible with our choice of C. At last, our only choice for the Green kernel is

$$\widetilde{G}(u,v) = \widetilde{F}(u,v)\widetilde{G}(v,v)\,, \quad u,v \in T\,.$$

Now we finally arrive at the only way how to define the transition probabilities, via (11.7):

$$p(u,v) = \frac{1}{\widetilde{G}(u,u)}\frac{\widetilde{F}(u,v)}{1-\widetilde{F}(u,v)\widetilde{F}(v,u)}. \tag{11.15}$$

Claim 1. $\mathcal{P}$ is stochastic.

Proof of Claim 1. Combining (11.15) with (11.14), we deduce that we have to verify that for every $u \in T$,

$$\sum_{v:v\sim u} \frac{\widetilde{F}(u,v)\big(1-\widetilde{F}(v,u)\big)}{1-\widetilde{F}(u,v)\widetilde{F}(v,u)} = 1\,. \tag{11.16}$$

If $u = o$, then by (11.12) this is just

$$\sum_{v:v\sim o} \mu(\partial T_v) = 1.$$

If $u \neq o$ then, again by (11.12), the left hand side of (11.16) is

$$\begin{aligned}\sum_{v:v^-=u} &\frac{\widetilde{F}(u,v)\big(1-\widetilde{F}(v,u)\big)}{1-\widetilde{F}(u,v)\widetilde{F}(v,u)} + \frac{\widetilde{F}(u,u^-)\big(1-\widetilde{F}(u^-,u)\big)}{1-\widetilde{F}(u,u^-)\widetilde{F}(u^-,u)} \\ &= \sum_{v:v^-=u} \frac{\mu(\partial T_v)}{\widetilde{F}(o,u)} + 1 - \frac{1-\widetilde{F}(u^-,u)}{1-\widetilde{F}(u,u^-)\widetilde{F}(u^-,u)} = 1.\end{aligned}$$

This proves Claim 1.

Claim 2. For any $u_0 \in T$, the function $\tilde{g}_{u_0}(u) = \widetilde{G}(u,u_0)$ satisfies $\mathcal{P}\tilde{g}_{u_0} = \tilde{g}_{u_0} - \mathbf{1}_{u_0}$.

Proof of Claim 2. First, we combine (11.14) with (11.15) to get

$$P\tilde{g}_{u_0}(u_0) = \sum_{v:v\sim u_0} p(u_0,v)\widetilde{F}(v,u_0)\widetilde{G}(u_0,u_0) = \sum_{v:v\sim u_0} \frac{\widetilde{F}(u_0,v)\widetilde{F}(v,u_0)}{1-\widetilde{F}(u_0,v)\widetilde{F}(v,u_0)} = \tilde{g}_{u_0}(u_0) - 1\,,$$

and Claim 2 is true at $u = u_0$. Second, for $u \neq u_0$, let w be the neighbour of u on $\pi(u,u_0)$. Then

$$\begin{aligned} P\tilde{g}_{u_0}(u) &= \sum_{v:v\sim u, v\neq w} p(u,v)\widetilde{F}(v,u)\widetilde{G}(u,u_0) + p(u,w)\widetilde{G}(w,u_0) \\ &= \underbrace{\sum_{v:v\sim u} \frac{\widetilde{F}(u,v)\widetilde{F}(v,u)}{1-\widetilde{F}(u,v)\widetilde{F}(v,u)}}_{\widetilde{G}(u,u)-1} \frac{\widetilde{G}(u,u_0)}{\widetilde{G}(u,u)} - p(u,w)\widetilde{F}(w,u)\widetilde{G}(u,u_0) + p(u,w)\widetilde{G}(w,u_0) \\ &= G(u,u_0)\left(1 - \frac{1}{\widetilde{G}(u,u)} - p(u,w)\widetilde{F}(w,u) + p(u,w)\frac{1}{\widetilde{F}(u,w)}\right) = \tilde{g}_{u_0}(u) \end{aligned}$$

since

$$p(u,w)/\widetilde{F}(u,w) - p(u,w)\widetilde{F}(w,u) = 1/\widetilde{G}(u,u)$$

by (11.15). This completes the proof of Claim 2.

Now we can conclude: the function $\tilde{g}_{u_0}$ is non-constant, positive and superharmonic. Therefore the random walk with transition matrix $\mathcal{P}$ given by 11.15 is transient and does posses a Green function $G(u,v)$. Furthermore, by the Riesz decomposition theorem, we have

$$\tilde{g}_{u_0} = Gf + h\,,$$

where h is a non-negative harmonic function and the charge f of the potential

$$Gf(u) = \sum_v G(u,v)f(v)$$

is $f = \tilde{g}_{u_0} - P\tilde{g}_{u_0} = \mathbf{1}_{u_0}$. That is,

$$\widetilde{G}(u,u_0) = G(u,u_0) + h(x) \quad \text{for all } u \in T.$$

Now let $x \in \partial T$ and $v = u_0 \wedge x$. If $u \in T_v$ then by our construction

$$\widetilde{G}(u,u_0) = \widetilde{G}(u,o)\frac{\widetilde{G}(v,u_0)}{\phi(v)}\phi(u) \to 0 \quad \text{as } u \to x\,.$$

Therefore $\widetilde{G}(\cdot, u_0)$ vanishes at infinity, and the same must hold for h. By the maximum principle, $h \equiv 0$.

We conclude that $\widetilde{G}(u,v) = G(u,v)$ for all $u, v \in T$. But then, by our construction, also $\widetilde{F}(u,v) = F(u,v)$, the "first hitting" kernel associated with $\mathcal{P}$. Comparing (11.12) with (10.10), we see that $\mu = \nu_o$. This completes the proof. ■

We conclude this section with some remarks on possible extensions.

Remark 11.6 In the present notes, we have restricted attention to *compact* ultra-metric spaces for two reasons. First, Kigami's approach [24] starting with a random walk on a rooted, locally finite tree T is restricted to that situation, because ∂T is compact.

On the other hand, our general approach in the present work is not restricted to compact spaces. In case of a non-compact, locally compact ultra-metric space without isolated points, one constructs the tree in the same way (the vertex set corresponds to the collection of all closed balls), but then the tree will have "its root at infinity", i.e., the ultra-metric space becomes $\partial^* T = \partial T \setminus \{\varpi\}$, where ϖ is a fixed reference end of T. In this situation, the predecessor v^- of a vertex v is the neighbour of v on $\pi(v, \varpi)$. In the definition 10.1 of an ultra-metric element, we then need besides monotonicity that ϕ tends to ∞ along $\pi(x, \varpi)$, while it has to tend to 0 along any geodesic going to $\partial^* T$. In this setting, the reference measure μ of a (ϕ, μ, σ)-process may have infinite mass: a Radon measure supported on the whole of $\partial^* T$.

In the non-compact case, however – and this is the second reason – there is not such an elegant and concise interpretation (analogous to the *boundary process*) of a (ϕ, μ, σ)-process in terms of a random walk. Nevertheless, in a very recent follow-up of [24], Kigami [25] studies a relation between random walks on a tree T and the non-compact ultrametric space $\partial^* T$. We intend to come back to this issue within the context of our approach.

Remark 11.7 Here, we have always assumed that the ultra-metric space has no isolated points, which for the tree means that $\deg^+ \geq 2$. Theme of [5] is the opposite situation, where all points are isolated, i.e., the space is discrete. In that case the ultrametric space also is the boundary of a tree, which does not consist of ends, but of terminal vertices, that is, vertices with only one neighbour.

From the point of view of the present section, the mixed situation works equally well. If we start with a locally compact ultra-metric space having both isolated and non-isolated points, we can construct the tree in the same way. The vertex set is the collection of all closed balls. The isolated points will then become *terminal vertices* of the tree, which have no neighbour besides the predecessor. All interior (non terminal) vertices will have forward degree ≥ 2.

In the compact case, the boundary ∂T of that tree should consist of the terminal vertices together with the space of ends. In the non-compact case, we will again have a reference end ϖ as in Remark 11.6.

The definition of an ultra-metric element remains the same, but we only need to define it on interior vertices. In this general setting, the construction of (ϕ, μ, σ) processes remains unchanged.

When the ultra-metric space is compact, even in presence of isolated points, the duality with random walks on the associated tree remains as explained here. The random walk should then be such that the terminal vertices are absorbing, and that the Green kernel tends to 0 at infinity. The Doob-Naïm formula extends readily to that setting.

Remark 11.8 Let us now consider the general (compact) situation when we start with a transient random walk on a locally finite, rooted tree T.

The limit distribution ν_o will in general not be supported by the whole of ∂T. The boundary process can of course still be constructed, see [24], but will evolve naturally on $\mathrm{supp}(\nu_o)$ only. Thus, we can consider our ultra-metric space to be just $\mathrm{supp}(\nu_o)$. The tree associated with this ultra-metric space will in general not be the tree we started with, nor its *transient skeleton* as defined in [47, (9.27)] (the subtree induced by o and all $v \in T \setminus \{o\}$ with $F(v, v^-) < 1$).

The reasons are twofold. First, the construction of the tree associated with $\mathrm{supp}(\nu_o)$ will never give back vertices with forward degree 1. Second, some end contained in $\mathrm{supp}(\nu_o)$ may be isolated within that set, while not being isolated in ∂T. But then this element will become a terminal vertex in the tree associated with the ultra-metric (sub)space $\mathrm{supp}(\nu_o)$. This occurs precisely when the transient skeleton has isolated ends.

Thus, one should work with a modified "reduced" tree plus random walk in order to maintain the duality between random walks and isotropic jump processes.

Remark 11.9 Given a transient random walk on the tree T, [24] also recovers an *intrinsic metric* of the boundary process on ∂T in terms of what is called an ultra-metric element in the present paper. This is of course $\phi(x) = G(x, o)$, denoted D_x in [24], where it is shown that for ν_o-almost every $\xi \in \partial T$, $D_x \to 0$ along the geodesic ray $\pi(o, \xi)$. This has the following Potential theoretic interpretation.

A point $x \in \partial T$ is called *regular for the Dirichlet problem,* if for every $\varphi \in C(\partial T)$, its Poisson transform h_φ satisfies

$$\lim_{u \to x} h_\varphi(u) = \varphi(x).$$

It is known from Cartwright, Soardi and Woess [13, Remark 2] that x is regular if and only if $\lim_{u \to x} G(u, o) = 0$ (as long as T has at least 2 ends), see also [47, Theorem 9.43]. By that Theorem 9.43, the set of regular points has ν_o-measure 1. That is, the Green kernel vanishes at ν_o-almost every boundary point.

Remark 11.10 In the proof of Theorem 11.5, we have reconstructed random walk transition probabilities from $C \cdot \phi(u) = G(u, o)$ and $\mu = \nu_o$.

A similar (a bit simpler) question was addressed by Vondraček [45]: how to reconstruct the transition probabilities from *all* limit distributions ν_u, $u \in T$, on the boundary. This, as well as our method, basically come from (10.10) and (11.7) + (11.8), which can be traced back to Cartier [11].

12 Fractional derivative on p-adic integers and the associated random walk

In conclusion we consider a specific example which unites the approaches of Sections 8-9 and Sections 10-11.

The p-adic fractional derivative on the group Z_p. Let $\mathbb{Z}_p \subset \mathbb{Q}_p$ be the group of p-adic integers. As a counterpart of the operator D^α we introduce the operator of fractional derivative defined on $\mathbb{Z}_p$. We denote this operator $\mathbb{D}^\alpha$. We show that $\mathbb{D}^\alpha$ is the Laplacian of an appropriate isotropic Markov semigroup. Then in the next subsection we construct a random walk associated in the sense of Sections 10-11 with the operator $\mathbb{D}^\alpha$.

Since $\mathbb{Z}_p$ is a compact Abelian group, its dual $\widehat{\mathbb{Z}_p}$ is a discrete Abelian group. It is known that the group $\widehat{\mathbb{Z}_p}$ can be identified with the group

$$Z(p^\infty) = \{p^{-n}m : 0 \le m < p^n, n = 1, 2, ...\}$$

equipped with the group operation addition of numbers mod 1. As sets (but not as groups) $Z(p^\infty) \subset \mathbb{Q}_p$, whence the function $\xi \mapsto \|\xi\|_p$ is well-defined on the group $Z(p^\infty)$.

Definition 12.1 The operator $\mathbb{D}^\alpha$ acting on the Abelian group $\mathbb{Z}_p$ is the L^2-multiplicator with symbol $\widehat{\mathbb{D}^\alpha}(\xi) = \|\xi\|_p^\alpha$, that is,

$$\widehat{\mathbb{D}^\alpha f}(\xi) = \|\xi\|_p^\alpha \widehat{f}(\xi), \quad \xi \in Z(p^\infty).$$

(Compare with the Definition 8.1 of the operator D^α.)

An immediate consequence of the Definition 12.1 is that the operator $\mathbb{D}^\alpha$ is a non-negative definite self-adjoint operator whose spectrum coincides with the range of the function

$$\xi \mapsto \|\xi\|_p^\alpha : Z(p^\infty) \to \mathbb{R}_+,$$

that is,

$$\mathrm{Spec}(\mathbb{D}^\alpha) = \{0, p^\alpha, p^{2\alpha}, ...\}.$$

The eigenspace $\mathcal{H}(\lambda)$ of the operator $\mathbb{D}^\alpha$ corresponding to the eigenvalue $\lambda = p^{k\alpha}$, $k \geq 1$, is spanned by the functions

$$f_k = \frac{1}{\mu_p(p^k\mathbb{Z}_p)}\mathbf{1}_{p^k\mathbb{Z}_p} - \frac{1}{\mu_p(p^{k-1}\mathbb{Z}_p)}\mathbf{1}_{p^{k-1}\mathbb{Z}_p}$$

and their shifts $f_k(\cdot + a)$ with any $a \in \mathbb{Z}_p/p^k\mathbb{Z}_p$.

Indeed, computing the Fourier transform of the function f_k,

$$\widehat{f_k}(\xi) = \mathbf{1}_{\{\|\xi\|_p \leq p^k\}} - \mathbf{1}_{\{\|\xi\|_p \leq p^{k-1}\}} = \mathbf{1}_{\{\|\xi\|_p = p^k\}},$$

we obtain

$$\widehat{\mathbb{D}^\alpha f_k}(\xi) = \|\xi\|_p^\alpha \widehat{f_k}(\xi) = p^{k\alpha}\widehat{f_k}(\xi).$$

The maximal number of linearly independent functions in the set $\{f_k(\cdot + a) : a \in \mathbb{Z}_p/p^k\mathbb{Z}_p\}$ is $p^{k-1}(p-1)$, whence

$$\dim \mathcal{H}(\lambda) = p^{k-1}(p-1).$$

All the above shows that $\mathbb{D}^\alpha$ coincides with the Laplacian (minus Markov generator) of some isotropic Markov semigroup $(\mathbb{P}_t^\alpha)_{t>0}$ defined on the ultra-metric measure space $(\mathbb{Z}_p, d_p, \mu_p)$ as constructed in (1.2) – (1.4) and studied in the previous sections. In particular, using the complete description of the set $\mathrm{Spec}(\mathbb{D}^\alpha)$ we compute the intrinsic distance, call it $d_{p,\alpha}(x,y)$,

$$d_{p,\alpha}(x,y) = \left(\frac{\|x-y\|_p}{p}\right)^\alpha.$$

It is now straightforward to compute the spectral distribution function $\mathbb{N}_\alpha(x,\tau) \equiv \mathbb{N}_\alpha(\tau)$ and then the jump-kernel $\mathbb{J}_\alpha(x,y) \equiv \mathbb{J}_\alpha(x-y)$ of the operator $\mathbb{D}^\alpha$. We claim that

$$\mathbb{J}_\alpha(x,y) = \frac{p^\alpha - 1}{1-p^{-\alpha-1}}\left(\frac{p^{-\alpha} - p^{-\alpha-1}}{1-p^{-\alpha}} + \frac{1}{\|x-y\|_p^{1+\alpha}}\right). \tag{12.1}$$

Recall for comparison that according to (8.3) the jump-kernel $J_\alpha(x,y)$ of the operator D^α is given by

$$J_\alpha(x,y) = \frac{p^\alpha - 1}{1-p^{-\alpha-1}}\frac{1}{\|x-y\|_p^{1+\alpha}}.$$

To prove (12.1), we compute $\mathbb{J}_\alpha(z)$. Let $\|z\|_p = p^{-l}$, then $d_{p,\alpha}(0,z) = p^{-(l+1)\alpha}$ and

$$\mathbb{J}_\alpha(z) = \int\limits_0^{1/d_\alpha(0,z)} \mathbb{N}_\alpha(\tau)d\tau = \int\limits_0^{p^{(l+1)\alpha}} \mathbb{N}_\alpha(\tau)d\tau.$$

The function $\mathbb{N}_\alpha(\tau)$ is a non-decreasing, left-continuous staircase function having jumps at the points $\tau_k = p^{k\alpha}$, $k = 1, 2, ...$, and taking values at these points $\mathbb{N}_\alpha(\tau_k) = p^{k-1}$, whence

$$\begin{aligned}\mathbb{J}_\alpha(z) &= 1 \cdot p^\alpha + p(p^{2\alpha} - p^\alpha) + p^2(p^{3\alpha} - p^{2\alpha}) + ... + p^l(p^{(l+1)\alpha} - p^{l\alpha}) \\ &= \frac{1 - p^{-1}}{1 - p^{-\alpha-1}} + \frac{p^\alpha - 1}{1 - p^{-\alpha-1}} p^{l(\alpha+1)} \\ &= \frac{p^\alpha - 1}{1 - p^{-\alpha-1}} \left(\frac{p^{-\alpha} - p^{-\alpha-1}}{1 - p^{-\alpha}} + \frac{1}{\|z\|_p^{1+\alpha}} \right)\end{aligned}$$

as desired.

Next we apply Theorem 7.2 and obtain

$$\mathbb{D}^\alpha f(x) = \int\limits_{\mathbb{Z}_p} (f(x) - f(y))\, \mathbb{J}_\alpha(x-y)\, d\mu_p(y) \tag{12.2}$$

and

$$(\mathbb{D}^\alpha f, f) = \frac{1}{2} \int\limits_{\mathbb{Z}_p \times \mathbb{Z}_p} (f(x) - f(y))^2\, \mathbb{J}_\alpha(x-y)\, d\mu_p(x)\, d\mu_p(y).$$

The equations (12.1)-(12.2) and (8.2) now yield the following result.

Corollary 12.2 *For any function f defined on $\mathbb{Z}_p \subset \mathbb{Q}_p$ we set $\widetilde{f} := f$ on $\mathbb{Z}_p$ and 0 otherwise. Then,*

$$f \in \mathsf{Dom}(\mathbb{D}^\alpha) \Longrightarrow \widetilde{f} \in \mathsf{Dom}(D^\alpha),$$

$$\mathbb{D}^\alpha f(x) = D^\alpha \widetilde{f}(x) \quad \text{and} \quad (\mathbb{D}^\alpha f, f) = \left(D^\alpha \widetilde{f}\, , \widetilde{f}\, \right) \tag{12.3}$$

whenever $x \in \mathbb{Z}_p$, $f \in \mathsf{Dom}(\mathbb{D}^\alpha)$ and $(\mathbf{1}, f) = 0$.

Nearest neighbour random walk on the tree T_p. As an illustration of Theorem 11.5 we construct a random walk on the tree $\mathbb{T}_p$ whose boundary process coincides with the isotropic process driven by the operator $C \cdot \mathbb{D}^\alpha$, where $C = p^{-\alpha}(1 - p^{-\alpha})$.

The Abelian group $\mathbb{Z}_p$ can be identified with the boundary of the tree $\mathbb{T}_p$ with root o where every vertex v has p forward neighbors. In our identification, $\mathbb{T}_p$ is the tree of balls of the ultra-metric space $(\mathbb{Z}_p, d_p)$ with root o corresponding to the whole of $\mathbb{Z}_p$ and the ultra-metric $d_p(x, y) = \|x - y\|_p$. We fix a constant $c \in (0, 1)$ and consider on $\mathbb{T}_p$ the nearest neighbour random walk with

$$p(v^-, v) = \begin{cases} 1/p & \text{if } v^- = o \\ c/p & \text{otherwise} \end{cases} \quad \text{and } p(v, v^-) = 1 - c.$$

Using [47, Thm. 1.38 and Prop. 9.3] one can compute precisely the Green function $G(v, o)$, the hitting probability $F(v, o)$ and other quantities associated with our random walk. In particular, choosing $c = (1 + p^{-\alpha})^{-1}$, we obtain

$$F(v, o) = p^{-\alpha|v|} \quad \text{and} \quad G(v, o) = \frac{p^{-\alpha|v|}}{1 - p^{-\alpha}}, \tag{12.4}$$

where $|v|$ is the graph distance from v to o. We see that the Green function vanishes at infinity, whence the random walk is Dirichlet regular.

The transition probabilities are invariant with respect to all automorphisms of the tree. Every such automorphism must fix o and every level of the tree. Let $\nu = \nu_o$ be the limit distribution on

$\partial\mathbb{T}_p$ of the random walk starting at o. Then also ν is invariant under the automorphism group of $\mathbb{T}_p$ (whose action extends to the boundary). In particular it is invariant under the action of $\mathbb{Z}_p$. Thus, under the identification of $\partial\mathbb{T}_p$ with $\mathbb{Z}_p$, we have that $\nu = \mu_p$, the normalized Haar measure of $\mathbb{Z}_p$.

We now look at the boundary process induced by our random walk as a jump process on $\mathbb{Z}_p$. By Theorem 11.1, the boundary process arises as an isotropic jump process with the reference measure μ_p. Let $\mathcal{L}$ be its Laplacian. By Corollary 11.2, the set $\mathrm{Spec}(\mathcal{L})$ coincides with the range of the function $v \mapsto 1/G(v,o), v \in \mathbb{T}_p$, plus $\{0\}$. In view of the above formula for $G(v,o)$ this means that

$$\mathrm{Spec}(\mathcal{L}) = \{0, (1-p^{-\alpha}), p^{\alpha}(1-p^{-\alpha}), p^{2\alpha}(1-p^{-\alpha}), ...\}.$$

Remember that

$$\mathrm{Spec}(\mathbb{D}^{\alpha}) = \{0, p^{\alpha}, p^{2\alpha}, ...\} = \frac{p^{\alpha}}{1-p^{-\alpha}}\ \mathrm{Spec}(\mathcal{L}).$$

Since both $\mathbb{D}^{\alpha}$ and $\mathcal{L}$ have the same orthonormal basis of eigenfunctions, we conclude that they are proportional, that is,

$$\mathbb{D}^{\alpha} = \frac{p^{\alpha}}{1-p^{-\alpha}}\ \mathcal{L}. \tag{12.5}$$

Thus, the boundary process $\{X_t\}_{t>0}$ associated with our random walk and the isotropic jump process $\{X_t^{\alpha}\}_{t>0}$ driven by the operator $\mathbb{D}^{\alpha}$ are related by the linear time change $X_{t/C} = X_t^{\alpha}$, where $C = p^{-\alpha}(1-p^{-\alpha})$.

The equation (12.5) implies that the jump kernels $\mathbb{J}_{\alpha}(x,y)$ and $\Theta_o(x,y)$ of the operators $\mathbb{D}^{\alpha}$ and $\mathcal{L}$ respectively are related by

$$\mathbb{J}_{\alpha}(x,y) = \frac{p^{\alpha}}{1-p^{-\alpha}}\ \Theta_o(x,y). \tag{12.6}$$

We conclude this section showing how to compute the Naïm kernel

$$\Theta_o(x,y) = \frac{1}{G(o,o)F(o,v)F(v,o)}, \quad \text{where } v = x \wedge y,$$

directly, using the data in (12.4). We do not have yet $F(o,v)$. We shall compute

$$N(v) := \frac{1}{F(o,v)F(v,o)}.$$

Since it depends only on the level k of v, we consider an arbitrary geodesic ray $[o = v_0, v_1, ...]$ and set up a linear recursion for $N(v_k)$. Denoting by w_1 an arbitrary neighbour of o different from v_1 and applying [47, Prop. 9.3(b)] and (12.4), we obtain

$$F(o,v_1) = \frac{1}{p} + \frac{p-1}{p}F(w_1,o)F(o,v_1) = \frac{1}{p} + \frac{(p-1)p^{-\alpha}}{p}F(o,v_1),$$

whence

$$F(o,v_1) = \frac{1}{p^{\alpha}\,(p^{\alpha}-p+1)}.$$

Thus, we get the initial values

$$N(v_0) = 1 \text{ and } N(v_1) = p^{\alpha} - p + 1.$$

Next, for $k \geq 1$, we let w_{k+1} be a forward neighbour of v_k different from v_{k+1}. Applying once again [47, Prop. 9.3(b)] and (12.4), we obtain

$$\begin{aligned} F(v_k, v_{k+1}) \quad &= \quad \frac{p^{\alpha}}{p(p^{\alpha}+1)} + \frac{(p-1)p^{\alpha}}{p(p^{\alpha}+1)}F(w_{k+1}, v_k)F(v_k, v_{k+1}) \\ &\quad + \frac{1}{p^{\alpha}+1}F(v_{k-1}, v_k)F(v_k, v_{k+1}). \end{aligned}$$

We insert the value $F(w_{k+1}, v_k) = p^{-\alpha}$ and divide by

$$F(o, v_{k+1}) = F(o, v_k)F(v_k, v_{k+1}) = F(o, v_{k-1})F(v_{k-1}, v_k)F(v_k, v_{k+1}).$$

Then we get

$$\frac{1}{F(o, v_k)} = \frac{p^\alpha}{p(p^\alpha+1)}\frac{1}{F(o, v_{k+1})} + \frac{p-1}{p(p^\alpha+1)}\frac{1}{F(o, v_k)} + \frac{1}{p^\alpha+1}\frac{1}{F(o, v_{k-1})}.$$

Now we multiply both sides with $1/F(v_k, o) = p^{\alpha k}$ and get

$$N(v_k) = \frac{1}{p(p^\alpha+1)}N(v_{k+1}) + \frac{p-1}{p(p^\alpha+1)}N(v_k) + \frac{p^\alpha}{p^\alpha+1}N(v_{k-1}).$$

This is a homogeneous second order linear recursion with constant coefficients. Its characteristic polynomial has roots 1 and $p^{\alpha+1}$. Therefore

$$N(v_k) = A + Bp^{(\alpha+1)k}.$$

Inserting the initial values we easily find the values of A and B. In order to get the Naïm kernel, we have to multiply by $1/G(o, o) = 1 - p^{-\alpha}$. Thus finally, we get

$$\begin{aligned}
\Theta_o(x, y) &= \frac{(1-p^{-\alpha})(p-1)}{p^{\alpha+1}-1} + \frac{(1-p^{-\alpha})(p^{\alpha+1}-p)}{p^{\alpha+1}-1}p^{(\alpha+1)k} \\
&= \frac{1-p^{-\alpha}}{p^\alpha}\frac{p^\alpha-1}{1-p^{-\alpha-1}}\left(\frac{p^{-\alpha}-p^{-\alpha-1}}{1-p^{-\alpha}} + \frac{1}{\|x-y\|_p^{\alpha+1}}\right) \\
&= \frac{1-p^{-\alpha}}{p^\alpha}\mathbb{J}_\alpha(x, y)
\end{aligned}$$

as desired.

References

[1] S. Albeverio and W. Karwowski, *A random walk on p-adics: generator and its spectrum*, Stochastic Processes and their Applications **53** (1994), 1–22.

[2] S. Albeverio and X. Zhao, *On the relation between different constructions of random walks on p-adics*, Markov Process. Related Fields **6** (2000), 239–255.

[3] A. Bendikov, B. Bobikau, and Ch. Pittet, *Some spectral and geometric aspects of countable groups*, Random Walks, Boundaries and Spectra, Progress in Probability, vol. **64**, Birkhäuser, Basel, 2011, pp. 227–234.

[4] ———, *Spectral properties of a class of random walks on locally finite groups*, Groups, Geometry and Dynamical Systems, in print (2013).

[5] A. Bendikov, A. Grigor'yan, and Ch. Pittet, *On a class of Markov semigroups on discrete ultra-metric spaces*, Potential Anal. **37** (2012), 125–169.

[6] A. Bendikov and L. Saloff-Coste, *Random walks driven by low moment measures*, Ann. Probab. **40** (2012), 2539–2588.

[7] ———, *Random walks on groups and discrete subordination*, Math. Nachr. **285** (2012), 580–605.

[8] Alexander Bendikov, *Lecture notes on Markov semigroups defined on totally disconnected sets*, Wrocław University, 2013.

[9] N. H. Bingham, C. M. Goldie, and J. L. Teugels, *Regular variation*, Encyclopedia of Mathematics and its Applications, vol. **27**, Cambridge University Press, Cambridge, 1989.

[10] S. Brofferio and W. Woess, *On transience of card shuffling*, Proc. Amer. Math. Soc. **129** (2001), 1513–1519.

[11] P. Cartier, *Fonctions harmoniques sur un arbre*, Symposia Math. **9** (1972), 203–270.

[12] D. I. Cartwright, *Random walks on direct sums of discrete groups*, J. Theoret. Probab. **1** (1988), 341–356.

[13] D. I. Cartwright, P. M. Soardi, and W. Woess, *Martin and end compactifications of non locally finite graphs*, Transactions Amer. Math. Soc. **338** (1993), 670–693.

[14] T. Coulhon, A. Grigor'yan, and Ch. Pittet, *A geometric approach to on-diagonal heat kernel low bounds on groups*, Ann. Inst. Fourier (Grenoble). **51** (2001), 1763–1827.

[15] D. A. Darling and P. Erdös, *On the recurrence of a certain chain*, Proc. Amer. Math. Soc. **13** (1962), 447–450.

[16] J. L. Doob, *Boundary properties for functions with finite dirichlet integrals*, Ann. Inst. Fourier (Grenoble) **12** (1962), 573–621.

[17] N. Fereig and S. A. Molchanov, *Random walks on abelian groups with an infinite number of generators*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. **5** (1978), 22–29.

[18] L. Flatto and J. Pitt, *Recurrence criteria for random walks on countable abelian groups*, Illinois J. Math. **18** (1974), 1–19.

[19] A. Georgakopoulos and V. A. Kaimanovich, *In preparation.*

[20] M. Gromov, *Asymptotic invariants of infinite groups*, Geometric group theory, Vol. 2 (Sussex, 1991), London Math. Soc. Lecture Note Ser., vol. **182**, Cambridge Univ. Press, Cambridge, 1993, pp. 1–295.

[21] E. Hewitt and K. A. Ross, *Abstract harmonic analysis, 1: Structure of topological groups, integration theory, group representations*, Grundlehren der math. Wiss., vol. **115**, Springer-Verlag, Berlin, 1963.

[22] M. A. Kasymdzhanova, *Recurrence of invariant markov chains on a class of abelian groups*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. **3** (1981), 3–7.

[23] H. Kesten and F. Spitzer, *Random walks on countably infinite abelian groups*, Acta. Math. **114** (1965), 237–265.

[24] J. Kigami, *Dirichlet forms and associated heat kernels on the cantor set induced by random walks on trees*, Adv. Math. **225** (2010), 2674–2730.

[25] ———, *Transitions on a noncompact cantor set and random walks on its defining tree*, Annales de l'Institut Henri Poincaré Prob.& Stat., in print (2013).

[26] A. N. Kochubei, *Pseudo-differential equations and stochastics over non-Archimedean fields*, Monographs and Textbooks in Pure and Applied Mathematics, vol. **244**, Marcel Dekker Inc., New York, 2001.

[27] G. F. Lawler, *Recurrence and transience for a card shuffling model*, Combin. Probab. Comput. **4** (1995), 133–142.

[28] M. Del Muto and A. Figà-Talamanca, *Diffusion on locally compact ultrametric spaces*, Expo. Math. **22** (2004), 197–211.

[29] ______, *Anisotropic diffusion on totally disconnected abelian groups*, Pacific J. Math. **225** (2006), 221–229.

[30] L. Naïm, *Sur le rôle de la frontière de r. s. martin dans la théorie du potentiel*, Ann. Inst. Fourier (Grenoble) **7** (1957), 183–281.

[31] J. Pearson and J. Bellissard, *Noncommutative riemannian geometry and diffusion on ultrametric cantor sets*, J. Noncommut. Geometry **3** (2009), 447–480.

[32] Ch. Pittet and L. Saloff-Coste, *On the stability of the behavior of random walks on groups*, J. Geom. Anal. **10** (2000), 713–737.

[33] ______, *On random walks on wreath products*, Ann. Probab. **30** (2002), 948–977.

[34] Christophe Pittet and Laurent Saloff-Coste, *Amenable groups, isoperimetric profiles and random walks*, Geometric group theory down under (Canberra, 1996), de Gruyter, Berlin, 1999, pp. 293–316.

[35] Luis Ribes and Pavel Zalesskii, *Profinite groups*, second ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, vol. **40**, Springer-Verlag, Berlin, 2010.

[36] L. Saloff-Coste, *Opérateurs pseudo-différentiels sur certains groupes totalement discontinus*, Studia Mathematica **83** (1986), 205–228.

[37] ______, *Probability on groups: random walks and invariant diffusions*, Notices Amer. Math. Soc. **48** (2001), 968–977.

[38] René L. Schilling, Renming Song, and Zoran Vondraček, *Bernstein functions*, second ed., de Gruyter Studies in Mathematics, vol. **37**, Walter de Gruyter & Co., Berlin, 2012.

[39] P. M. Soardi, *Potential theory on infinite networks*, Lecture Notes in Mathematics, vol. **1590**, Springer, Berlin, 1994.

[40] M. H. Taibleson, *Fourier analysis on local fields*, Princeton Univ. Press, 1975.

[41] N. Th. Varopoulos, L. Saloff-Coste, and T. Coulhon, *Analysis and geometry on groups*, Cambridge Tracts in Mathematics, vol. 100, Cambridge Univ. Press, Cambridge, 1992.

[42] V. S. Vladimirov, *Generalized functions over the field of p-adic numbers*, Uspekhi Mat. Nauk **43** (1988), 17–53, 239.

[43] V. S. Vladimirov and I. V. Volovich, *p-adic Schrödinger-type equation*, Lett. Math. Phys. **18** (1989), 43–53.

[44] V. S. Vladimirov, I. V. Volovich, and E. I. Zelenov, *p-adic analysis and mathematical physics*, Series on Soviet and East European Mathematics, vol. **1**, World Scientific Publishing Co., Inc., River Edge, NY, 1994.

[45] Z. Vondraček, *A characterization of markov chains on infinite graphs by limiting distributions*, Arch. Math. (Basel) **65** (1995), 449–460.

[46] Wolfgang Woess, *Random walks on infinite graphs and groups*, Cambridge Tracts in Mathematics, vol. **138**, Cambridge University Press, Cambridge, 2000.

[47] ______, *Denumerable Markov chains*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2009, Generating functions, boundary theory, random walks on trees.

[48] M. Yamasaki, *Discrete potentials on an infinite network*, Mem. Fac. Sci., Shimane Univ. **13** (1979), 31–44.

A. Bendikov: Institute of Mathematics, Wroclaw University
Pl. Grundwaldzki 2/4, 50-384 Wrocław, Poland
email: bendikov@math.uni.wroc.pl

A. Grigoryan: Department of Mathematics, University of Bielefeld,
33501 Bielefeld, Germany
email: grigor@math.uni-bielefeld.de

Ch. Pittet: CMI, Université d'Aix-Marseille I, 39 rue Joliot-Curie,
13453 Marseille Cedex 13, France
email: pittet@cmi.univ-mrs.fr

W. Woess: Institut für Mathematische Strukturtheorie, Technische Universität Graz
Steyrergasse 30, A-8010 Graz, Austria
email: woess@TUGraz.at